

2018

Smoothness of definite unitary eigenvarieties at critical points

John Bergdall

Bryn Mawr College, jbergdall@brynmawr.edu

[Let us know how access to this document benefits you.](#)

Follow this and additional works at: https://repository.brynmawr.edu/math_pubs

 Part of the [Mathematics Commons](#)

Custom Citation

Bergdall, John. "Smoothness of definite unitary eigenvarieties at critical points." *Journal für die reine und angewandte Mathematik*, forthcoming (published online 2018).

This paper is posted at Scholarship, Research, and Creative Work at Bryn Mawr College. https://repository.brynmawr.edu/math_pubs/21

For more information, please contact repository@brynmawr.edu.

SMOOTHNESS OF DEFINITE UNITARY EIGENVARIETIES AT CRITICAL POINTS

JOHN BERGDALL

ABSTRACT. We compute an upper bound for the dimension of the tangent spaces at classical points of certain eigenvarieties associated with definite unitary groups, especially including the so-called critically refined cases. Our bound is given in terms of “critical types” and when our bound is minimized it matches the dimension of the eigenvariety. In those cases, which we explicitly determine, the eigenvariety is necessarily smooth and our proof also shows that the completed local ring on the eigenvariety is naturally a certain universal Galois deformation ring.

1. INTRODUCTION

Let p be a prime number. Eigenvarieties, referring to p -adic families of automorphic forms, first appeared thirty years ago in the work of Hida on p -ordinary elliptic cuspidal eigenforms. Coleman and Mazur, and Buzzard, removed the ordinary condition in the final decade of the 20th century, constructing p -adic families passing through all finite slope cuspidal eigenforms. A number of authors have since given constructions of eigenvarieties in a wide range of situations. Despite the growing interest in the subject, basic geometric properties of eigenvarieties remain occluded.¹

The goal of this article is to give upper bounds for the dimensions of the tangent spaces at classical points on eigenvarieties associated with definite unitary groups. These bounds are given in terms of *critical types* of triangulations of local Galois representations at the p -adic places. The more critical a point is, the larger the upper bound.

We also give an exact condition for when our bound is minimized. In that case, our bound equals the dimension of the corresponding eigenvariety and we get a smoothness statement as well. For the rest of this introduction we set notation and state the main theorem.

1.1. Critical types of p -refined automorphic representations. Let F/F^+ be a CM extension of number fields such that each p -adic place of F^+ splits in F . We will use \mathbf{G} to denote a rank n unitary group associated to this extension. We assume $\mathbf{G} \times_{F^+} F \simeq \mathrm{GL}_{n/F}$ and that \mathbf{G} is compact at infinity. In the body of the text, we will also make further technical assumptions that we omit now (see Section 4). If v is a p -adic place of F^+ then the choice of a place $\tilde{v} \mid v$ in F defines an isomorphism $\mathbf{G}(F_v^+) \simeq \mathrm{GL}_n(F_{\tilde{v}}) = \mathrm{GL}_n(F_v^+)$. Thus, the local components of automorphic representations for \mathbf{G} at p -adic places are irreducible smooth representations of $\mathrm{GL}_n(F_v^+)$.

Associated with an automorphic representation π for \mathbf{G} is a p -adic Galois representation

$$\rho_\pi : G_F \rightarrow \mathrm{GL}_n(\overline{\mathbf{Q}}_p).$$

Date: October 19, 2017.

2000 Mathematics Subject Classification. 11F33 (11F80, 11F55).

¹For example: it is not known if the tame level 1, 2-adic eigencurve has finitely many connected components.

The representation ρ_π is geometric in the sense of Fontaine and Mazur: ρ_π is unramified except at finitely many places and it is de Rham at the p -adic places.

We now fix an automorphic representation π of \mathbf{G} lying in the unramified principal series at the p -adic places $v \mid p$ in F^+ . For such v , we also fix the choice of $\tilde{v} \mid v$ in F and denote by $\rho_{\pi, \tilde{v}}$ its restriction to a decomposition group at \tilde{v} . Since the local component π_v at a place $v \mid p$ is an unramified principal series, the representation $\rho_{\pi, \tilde{v}}$ is crystalline. Note that $F_{\tilde{v}}^- = F_v^+$ and, to emphasize that the choice of \tilde{v} does not matter substantially, we write F_v^+ for this p -adic field. Finally, we choose a finite extension L/\mathbf{Q}_p , inside a fixed algebraic closure $\overline{\mathbf{Q}_p}$, such that the image of ρ_π is contained in $\mathrm{GL}_n(L)$, and we assume that L contains the image of any embedding $\tau : F_v^+ \hookrightarrow \overline{\mathbf{Q}_p}$ for each v .

Along with π and the distinguished places \tilde{v} , we also fix a collection of triangulations $(P_{\tilde{v}, \bullet})_{\tilde{v}}$. Specifically, we consider the $(\varphi, \Gamma_{F_v^+})$ -module $D_{\pi, \tilde{v}} := D_{\mathrm{rig}}^\dagger(\rho_{\pi, \tilde{v}})$ over the Robba ring $\mathcal{R}_{F_v^+, L}$ (see Section 2), and the triangulation $P_{\tilde{v}, \bullet}$ is a full filtration

$$P_{\tilde{v}, \bullet} : (0) = P_{\tilde{v}, 0} \subsetneq P_{\tilde{v}, 1} \subsetneq \cdots \subsetneq P_{\tilde{v}, n-1} \subsetneq P_{\tilde{v}, n} = D_{\pi, \tilde{v}}$$

of saturated $(\varphi, \Gamma_{F_v^+})$ -submodules. We refer to π together with the choice of triangulations as a p -refined automorphic representation (the terminology goes back to [36]).

The graded pieces of the triangulation $P_{\tilde{v}, \bullet}$ have rank one and so $P_{\tilde{v}, \bullet}$ defines an ordered tuple $(\delta_{\tilde{v}, 1}, \dots, \delta_{\tilde{v}, n})$ of continuous characters $\delta_{\tilde{v}, i} : (F_v^+)^\times \rightarrow L^\times$ as in [30, Section 6.2] (see also Section 2.1). For each embedding $\tau : F_v^+ \hookrightarrow L$ there exists an integer $s_{i, \tilde{v}, \tau}$ (the τ -Hodge–Tate weight of $\delta_{\tilde{v}, i}$) such that $\delta_{\tilde{v}, i}(z) = \prod_\tau \tau(z)^{-s_{i, \tilde{v}, \tau}}$ for all z in $\mathcal{O}_{F_v^+}^\times$. (Our normalization gives the identity character Hodge–Tate weight -1 at each embedding τ .)

On the other hand, for each embedding $\tau : F_v^+ \hookrightarrow L$, we also have the (necessarily distinct) Hodge–Tate weights $h_{1, \tilde{v}, \tau} < h_{2, \tilde{v}, \tau} < \cdots < h_{n, \tilde{v}, \tau}$ of $D_{\pi, \tilde{v}}$. For fixed \tilde{v} and τ , the two sets $\{s_{i, \tilde{v}, \tau} : i = 1, \dots, n\}$ and $\{h_{i, \tilde{v}, \tau} : i = 1, \dots, n\}$ are equal, so we denote by $\sigma_{\tilde{v}, \tau}$ the unique permutation defined by $s_{i, \tilde{v}, \tau} = h_{\sigma_{\tilde{v}, \tau}(i), \tilde{v}, \tau}$.²

Definition 1.1. *The critical type of the triangulation $P_{\tilde{v}, \bullet}$ of $D_{\pi, \tilde{v}}$ is the collection of permutations $(\sigma_{\tilde{v}, \tau})_\tau$. The triangulation $P_{\tilde{v}, \bullet}$ is called non-critical if $\sigma_{\tilde{v}, \tau} = \mathrm{id}$ for each τ .*

The non-critical case is the most common. For instance, a p -refined automorphic representation of non-critical slope is non-critical (see [5, Remark 2.4.6(ii)]) and having non-critical slope is generic on an eigenvariety. Nevertheless, interesting arithmetic phenomena occur in critical situations (see [4, Theorem 2] for example) and it seems less difficult for a triangulation to be critical as $n \rightarrow \infty$. For contrast, if $n = 2$ and $F_v^+ = \mathbf{Q}_p$ then a critical triangulation at \tilde{v} exists if and only if $\rho_{\pi, \tilde{v}}$ is abelian.

1.2. Main result. An eigenvariety p -adically interpolates p -refined automorphic representations $(\pi, (P_{\tilde{v}, \bullet}))$. We refer to [20, 24] and Section 4 for details. Here, we fix a *minimal* eigenvariety X containing the pair $x := (\pi, (P_{\tilde{v}, \bullet}))$. Thus, X is a rigid analytic space over \mathbf{Q}_p , equidimensional of dimension $(F^+ : \mathbf{Q}) \cdot n$. It implicitly depends on the choice of a tame level; second, the minimal condition essentially means that the point x is not lying at the intersection of two eigenvarieties obtained from smaller tame levels. Our main theorem is a bound on the dimension of the Zariski tangent space $T_{X, x}$ of X at x in terms of critical types. To state it, we need two notations.

²It may make sense to replace $\sigma_{\tilde{v}, \tau}$ by its inverse (for formulae in representation theory to work out cleaner). The results below (Theorems 1.2 and 1.5) do not depend on this choice.

The representation ρ_π is conjugate self-dual, up to a twist, and if ρ_π is absolutely irreducible then the natural action of G_F on the adjoint representation $\text{ad } \rho_\pi$ extends to an action of the absolute Galois group G_{F^+} (see Section 4.3). We denote by $H_f^1(G_{F^+}, \text{ad } \rho_\pi)$ the corresponding Bloch–Kato Selmer group [13].

If σ is a permutation of $\{1, \dots, n\}$, we let $\ell(\sigma) = \{(i, j) : i < j \text{ and } \sigma(i) > \sigma(j)\}$. This is also the length of a minimal expression of σ as a product of simple transpositions. (A simple transposition is a transposition interchanging two consecutive integers i and $i + 1$.) We also write $c(\sigma)$ for the number of orbits of the group generated by σ acting on $\{1, \dots, n\}$. For example, $\ell(\text{id}) = 0$ and $c(\text{id}) = n$.

In the next theorem, we refer to Definition 3.4 for the notion of a regular generic triangulation. For an idea, “regular” is a simplicity condition on crystalline eigenvalues.

Theorem 1.2 (Theorem 4.7). *Suppose that π is an automorphic representation of \mathbf{G} which is unramified at each p -adic place and $(P_{\tilde{v}, \bullet})$ is a collection of regular generic triangulations for ρ_π at the p -adic places. Assume that ρ_π is irreducible and X is the minimal eigenvariety containing the point $x = (\pi, (P_{\tilde{v}, \bullet}))$. Then,*

$$\dim T_{X,x} \leq \dim H_f^1(G_{F^+}, \text{ad } \rho_\pi) + \sum_{\tilde{v}, \tau} \ell(\sigma_{\tilde{v}, \tau}) + c(\sigma_{\tilde{v}, \tau}),$$

where $(\sigma_{\tilde{v}, \tau})_{\tilde{v}, \tau}$ are the critical types of the triangulations at x .

We also have the following direct corollary.

Corollary 1.3 (Corollary 4.8). *With the notation and assumptions of Theorem 1.2, also assume that $H_f^1(G_{F^+}, \text{ad } \rho_\pi) = (0)$ and each $\sigma_{\tilde{v}, \tau}$ is a product of distinct simple transpositions. Then, X is smooth at the point x .*

The Selmer group is conjectured to vanish;³ it has been proven in many cases, under hypotheses inherent to the methods of Taylors–Wiles and Kisin (see [1] and [17, Section 4]). Denote by $\bar{\rho}_\pi^{\text{ss}}$ the semi-simplification of any mod p reduction of ρ_π . As a concrete version of Corollary 1.3, the main theorem of [1] implies:

Corollary 1.4. *With the notation and assumptions of Theorem 1.2, assume that each $\sigma_{\tilde{v}, \tau}$ is a product of distinct simple transpositions, $\zeta_p \notin F$, and $\bar{\rho}_\pi^{\text{ss}}(G_{F(\zeta_p)})$ is adequate.⁴ Then X is smooth at x .*

The main corollary is deduced from the theorem as follows. A short computation shows that the contribution of the critical types in Theorem 1.2 is minimized, and equal to $(F^+ : \mathbf{Q}) \cdot n$, exactly when each critical type is a product of distinct simple transpositions. Since X is equidimensional of dimension $(F^+ : \mathbf{Q}) \cdot n$, this means that X is regular, and thus smooth, at x in the situation of Corollary 1.3. The proof also gives an “ $R = \mathbf{T}$ ” theorem which we will partially explain in Section 1.3 below (see Corollary 4.8 for a precise statement).

Examples constructed in [3] by Bellaïche show that the irreducibility of ρ_π is important in Theorem 1.2 and Corollary 1.3. Regarding optimality, Breuil, Hellmann and Schraen have shown that Corollary 1.3 is optimal in that X is singular once one of its critical types is *not* a product of distinct simple transpositions. See [15, Theorem 1.2] and compare with the

³In fact, the larger space $H_f^1(G_F, \text{ad } \rho_\pi)$ is conjectured to vanish.

⁴The definition of adequate is taken from [1, Definition 3.1.1], which generalizes [44, Definition 2.3].

earlier result [16, Corollary 5.18]. We expand on these notes following Corollary 4.8 in the text.

1.3. Sketch of proof. The proof of Theorem 1.2 follows a well-known strategy: we compare a ring of Hecke operators to a universal deformation ring for a Galois representation. Our ring of Hecke operators is the completion $\widehat{\mathcal{O}}_{X,x}^{\text{rig}}$ of the rigid analytic local ring of the eigenvariety X at the point x , and the deformation ring, denoted $R_{\rho_\pi}^{\text{Ref,min}}$, is a deformation ring for ρ_π .

The deformations parameterized by $R_{\rho_\pi}^{\text{Ref,min}}$ are *weakly-refined* at p -adic places and *minimally ramified* (or, unramified in the sense of Bloch and Kato) at the places away from p . The weakly-refined condition depends on a triangulation (which is suppressed in the notation). When the triangulation is non-critical, the weakly-refined deformations are the same as the *trianguline* deformations studied in [5, Chapter 2].

The interpolation of crystalline periods over eigenvarieties [31, 34] and the minimality of our eigenvariety implies that there is a natural surjective map $R_{\rho_\pi}^{\text{Ref,min}} \twoheadrightarrow \widehat{\mathcal{O}}_{X,x}^{\text{rig}}$. In this way, a bound for the tangent space $T_{X,x}$ is obtained from any bound for the Zariski tangent space $\mathfrak{t}_{\rho_\pi}^{\text{Ref,min}}$ of the deformation ring $R_{\rho_\pi}^{\text{Ref,min}}$.

Tangent spaces of deformation rings are computed using Galois cohomology. Following an idea of Bellaïche and Chenevier in the non-critical case, we observe that the global tangent space $\mathfrak{t}_{\rho_\pi}^{\text{Ref,min}}$ is naturally equipped with restriction maps to the tangent spaces $\mathfrak{t}_{\rho_\pi,\tilde{v}}^{\text{Ref}}$ of the weakly refined deformation problem at the p -adic places (see Section 3.2). One has a natural exact sequence

$$(1) \quad 0 \rightarrow H_f^1(G_{F^+}, \text{ad } \rho_\pi) \rightarrow \mathfrak{t}_{\rho_\pi}^{\text{Ref,min}} \rightarrow \bigoplus_{v|p} \mathfrak{t}_{\rho_\pi,\tilde{v}}^{\text{Ref}} / H_f^1(G_{F_{\tilde{v}}}, \text{ad } \rho_\pi, \tilde{v}).$$

where $H_f^1(G_{F_{\tilde{v}}}, \text{ad } \rho_\pi, \tilde{v})$ is the local Bloch-Kato Selmer group parameterizing infinitesimal crystalline deformations of ρ_π, \tilde{v} .

Our main technical result is a bound on the third term in the sequence (1). The following is a purely local theorem, but we state it here in the global context where it is applied.

Theorem 1.5 (Theorem 3.16). *For each $v \mid p$,*

$$\dim \mathfrak{t}_{\rho_\pi,\tilde{v}}^{\text{Ref}} / H_f^1(G_{F_{\tilde{v}}}, \text{ad } \rho_\pi, \tilde{v}) \leq \sum_{\tau: F_v^+ \rightarrow L} \ell(\sigma_{\tilde{v},\tau}) + c(\sigma_{\tilde{v},\tau}).$$

Combining the bound in Theorem 1.5 with the sequence (1) and the preceding paragraphs, we get the bound in Theorem 1.2.

In the non-critical case, Theorem 1.5 is proven in [5] by computing the dimension of a trianguline deformation ring. The key point in our generalization is carefully measuring, in terms of the critical type, how far weakly-refined deformations are from being trianguline. For that, we separately study (I) weakly-refined deformations with constant Hodge–Tate weights and (II) the variation of Hodge–Tate weights in weakly-refined deformations.

Versions of the above results were obtained in low-dimensional cases by the author in his Ph.D. thesis [8] and later in unpublished notes. The explicit goal was to prove the smoothness part of Corollary 1.3, but only the second half of the computation, referring to (II), was well understood (see [9, Section 7] for example).

The condition of the critical types being products of distinct simple transpositions in Corollary 1.3 did not occur to the author until hearing in lectures at the Centre International de Rencontres Mathématiques (Luminy) in 2015 that a similar local result was proven by Breuil, Hellmann and Schraen [16]. Once the condition was noticed, the statement of Theorem 1.5 and its proof were obtained independently. Compare with [16, Section 4]. The applications to eigenvarieties in this paper and in [16] are different.

1.4. Organization. We give a brief reminder on (φ, Γ) -modules in Section 2. In Section 3 we define the weakly-refined deformations and prove Theorem 1.5. Section 4 is dedicated to the proofs of Theorem 1.2 and Corollary 1.3.

1.5. Notations and conventions. We fix an algebraic closure $\overline{\mathbf{Q}}_p$ and an isomorphism $\overline{\mathbf{Q}}_p \simeq \mathbf{C}$ which is used implicitly throughout. We assume that L is a finite extension of \mathbf{Q}_p contained in $\overline{\mathbf{Q}}_p$, and we allow L to change so as to contain the image of any embedding of a p -adic field into $\overline{\mathbf{Q}}_p$.

Suppose ℓ is a prime (possibly $\ell = p$) and K/\mathbf{Q}_ℓ is a finite extension. We write K_0 for the maximal unramified subextension of K and ℓ^{f_K} for the number of elements in the residue field of K . If $\rho : G_K \rightarrow \mathrm{GL}_n(\overline{\mathbf{Q}}_p)$ is a continuous representation which is potentially semi-stable (this is automatic if $\ell \neq p$) then we write $\mathrm{WD}(\rho)$ for the corresponding Weil–Deligne representation over $\overline{\mathbf{Q}}_p$ (see [42, Theorem 4.2.1] if $\ell \neq p$ and [25] if $\ell = p$). As an example, if $\ell = p$ and ρ is crystalline then $\mathrm{WD}(\rho)$ is unramified and the eigenvalues of a geometric Frobenius element are the eigenvalues of the crystalline Frobenius φ^{f_K} acting on $D_{\mathrm{crys}}(\rho)$, counted with multiplicity.

Let $\mathrm{rec}_K : K^\times \rightarrow G_K^{\mathrm{ab}}$ be the local Artin reciprocity map, normalized so that the image of a uniformizer corresponds to a geometric Frobenius element. If π is an irreducible smooth representation of $\mathrm{GL}_n(K)$ on a $\overline{\mathbf{Q}}_p$ -vector space we denote by $\mathrm{rec}(\pi)$ the n -dimensional Frobenius semi-simple Weil–Deligne representation over $\overline{\mathbf{Q}}_p$ given by the local Langlands correspondence [26]. We normalize the correspondence as follows. Let $T(K) \subset \mathrm{GL}_n(K)$ be the diagonal torus and $B(K) \subset \mathrm{GL}_n(K)$ be the upper triangular subgroup. If $\chi = \chi_1 \otimes \cdots \otimes \chi_n$ is a character of $T(K)$ then $\mathrm{rec}(\pi(\chi)) = \chi_1 \circ \mathrm{rec}_K^{-1} \oplus \cdots \oplus \chi_n \circ \mathrm{rec}_K^{-1}$, where $\pi(\chi)$ is the unique irreducible unramified subquotient of the smooth, non-normalized, induction $\mathrm{Ind}_{B(K)}^{\mathrm{GL}_n(K)}(\delta_{B(K)}^{1/2} \cdot \chi)$. Here, $\delta_{B(K)} = |-|_K^{n-1} \otimes \cdots \otimes |-|_K^{1-n}$ is the modulus character of $B(K)$.

If $\ell = p$, and $\delta : \mathcal{O}_K^\times \rightarrow L^\times$ is a continuous character then for each embedding $\tau : K \hookrightarrow L$ we write $\mathrm{HT}_\tau(\delta)$ for the τ -Hodge–Sen–Tate weight of δ , which is the negative of the weight defined in [30, Definition 6.1.6].

If $n \geq 1$ we let S_n denote the group of permutations on $\{1, \dots, n\}$. If $\sigma \in S_n$ we write $\ell(\sigma) = \{(i, j) : i < j \text{ and } \sigma(i) > \sigma(j)\}$ for its length and $c(\sigma)$ for the number of orbits in $\{1, \dots, n\}$ under the action of the group generated by σ .

1.6. Acknowledgements. The author thanks Christophe Breuil, David Hansen, Eugen Hellmann and Benjamin Schraen for helpful discussions and comments. The author also thanks the Centre International de Rencontres Mathématiques for hospitality in June 2015. Many ideas in this article were developed when the author was a graduate student at Brandeis University. Thanks are duly given to Joël Bellaïche for encouragement, numerous insightful conversations and comments on an early draft of this paper. The author was partially supported by NSF award DMS-1402005.

2. REMINDER ON (φ, Γ) -MODULES

2.1. (φ, Γ_K) -modules and triangulations. Let K/\mathbf{Q}_p be a finite extension. We denote by \mathcal{R}_K the Robba ring defined over K , i.e. the ring of series $f = \sum a_i T^i$ defined over the maximal absolutely unramified extension of K_∞ and which converge on an annulus $r(f) < |T| < 1$ (see [30, Section 2]). Here K_∞ is the field obtained from by adjoining all the p -power roots of unity to K . If A is an affinoid \mathbf{Q}_p -algebra then we define $\mathcal{R}_{K,A} := \mathcal{R}_K \widehat{\otimes}_{\mathbf{Q}_p} A$.

Recall that if D is a finite free module over a commutative ring R then we say a submodule $P \subset D$ is saturated if D/P is projective as an R -module. If L/\mathbf{Q}_p is a finite extension then $\mathcal{R}_{K,L}$ is an adequate Bézout domain [10, Proposition 4.12] and so, for $R = \mathcal{R}_{K,L}$, projective may be replaced for free. The following lemma is [5, Lemma 2.2.3] when $K = \mathbf{Q}_p$. The proof is no different for general K , so we omit it.

Lemma 2.1. *Let A be a local Artin L -algebra with residue field L and maximal ideal \mathfrak{m}_A . Suppose that D is a finite free $\mathcal{R}_{K,A}$ -module which contains a rank one free submodule $P \subset D$. If $P/\mathfrak{m}_A P \subset D/\mathfrak{m}_A D$ is saturated as an $\mathcal{R}_{K,L}$ -module then P is saturated in D as well.*

We equip $\mathcal{R}_{K,L}$ with its natural commuting actions of the Frobenius operator φ and the group $\Gamma_K = \text{Gal}(K_\infty/K)$ (see [30, Definition 2.2.2]). A (φ, Γ_K) -module D over $\mathcal{R}_{K,L}$ is a finite free $\mathcal{R}_{K,L}$ -module D equipped with commuting $\mathcal{R}_{K,L}$ -semilinear actions of an operator φ and the group Γ_K , such that $\varphi(D)$ generates D as an $\mathcal{R}_{K,L}$ -module. For coefficients more general than L (e.g. Artin algebras) see [5, Chapter 2] or [30, Section 2]. The rank of D is the rank of the underlying $\mathcal{R}_{K,L}$ -module. We write D^\vee for the dual (φ, Γ_K) -module.

There is a functor $\rho \mapsto D_{\text{rig}}^\dagger(\rho)$ which defines a fully faithful embedding

$$\{\text{continuous representations } \rho : G_K \rightarrow \text{GL}_n(L)\} \hookrightarrow \{\text{rank } n \text{ } (\varphi, \Gamma_K)\text{-modules over } \mathcal{R}_{K,L}\}.$$

Its essential image is the so-called étale (φ, Γ_K) -modules characterized using the theory of slope filtrations [28, Theorem 6.10]. Crucially, $D_{\text{rig}}^\dagger(\rho)$ may contain non-étale (φ, Γ_K) -submodules even if ρ is irreducible.

Rank one (φ, Γ_K) -modules over $\mathcal{R}_{K,L}$ are classified by continuous characters $\delta : K^\times \rightarrow L^\times$. We write $\mathcal{R}_{K,L}(\delta)$ for the (φ, Γ_K) -module corresponding to δ by [30, Construction 6.2.4]. If D is a (φ, Γ_K) -module over $\mathcal{R}_{K,L}$ then we write $D(\delta) := D \otimes_{\mathcal{R}_{K,L}} \mathcal{R}_{K,L}(\delta)$ for the “twist” of D by δ .

Important constructions in the theory of Galois representations extend to the category of (φ, Γ_K) -modules. For example, a (φ, Γ_K) -module has Galois cohomology $H^\bullet(D)$ concentrated in degree at most two ([27, 33]). If $\delta : K^\times \rightarrow L^\times$ is a continuous character we write $H^\bullet(\delta)$ in lieu of $H^\bullet(\mathcal{R}_{K,L}(\delta))$. We also have Fontaine’s notions of de Rham, crystalline, etc. for (φ, Γ_K) -modules. For example, $D_{\text{crys}}(D) = D[1/t]^{\Gamma_K}$ where $t \in \mathcal{R}_{\mathbf{Q}_p}$ is “Fontaine’s p -adic $2\pi i$ ”. (See [10, 11] for details.)

A triangulation of a (φ, Γ_K) -module D over $\mathcal{R}_{K,L}$ is a filtration

$$P_\bullet : 0 = P_0 \subsetneq P_1 \subsetneq \cdots \subsetneq P_{n-1} \subsetneq P_n = D$$

of D by saturated (φ, Γ_K) -submodules. The parameter of P_\bullet is the ordered tuple $(\delta_1, \dots, \delta_n)$ of continuous characters δ_j such that $P_j/P_{j-1} = \mathcal{R}_{K,L}(\delta_j)$.

Now let D be a crystalline (φ, Γ_K) -module over $\mathcal{R}_{K,L}$. Thus $D_{\text{crys}}(D)$ is a finite free $K_0 \otimes_{\mathbf{Q}_p} L$ -module equipped with a K_0 -semilinear (but L -linear) operator φ and $D_{\text{crys}}(D)_K := D_{\text{crys}}(D) \otimes_{K_0} K$ is equipped with a decreasing, exhaustive and separated filtration Fil^\bullet by $K \otimes_{\mathbf{Q}_p} L$ -submodules (the Hodge filtration). The operator φ^{f_K} is K_0 -linear and we refer to its

eigenvalues as the crystalline eigenvalues of D . Once L is sufficiently large, every crystalline eigenvalue lies in $L^\times \subset (K_0 \otimes_{\mathbf{Q}_p} L)^\times$ (compare with the proof of Lemma 3.3).

If P_\bullet is a triangulation of a crystalline (φ, Γ_K) -module D then $D_{\text{crys}}(P_j) \subset D_{\text{crys}}(D)$ is a filtered φ -submodule of rank j . Thus, a triangulation defines an ordering (ϕ_1, \dots, ϕ_n) of the crystalline eigenvalues by declaring the first j eigenvalues appear in $D_{\text{crys}}(P_j)$. If D has distinct crystalline eigenvalues this defines a bijection (see [11])

$$(2) \quad \{\text{triangulations of } D\} \longleftrightarrow \{\text{orderings of crystalline eigenvalues for } D\}.$$

Let us briefly recall the ‘‘matching’’ of weights in this bijection. If $\tau : K \hookrightarrow L$ is an embedding, the filtration Fil^\bullet on $D_{\text{crys}}(D)_K$ equips $D_{\text{crys}}(D)_K \otimes_{K,\tau} L$ with an exhaustive and separated filtration by L -subspaces whose jumps are the τ -Hodge–Tate weights $h_{1,\tau} \leq h_{2,\tau} \leq \dots \leq h_{n,\tau}$. Given an ordering (ϕ_1, \dots, ϕ_n) and an embedding τ we write $(s_{1,\tau}, \dots, s_{n,\tau})$ for the re-ordering of $\{h_{i,\tau}\}$ such that the induced filtration on $\sum_{i=1}^j D_{\text{crys}}(D)^{\varphi^i K = \phi_i} \otimes_{K_0,\tau} L$ has weights $\{s_{1,\tau}, \dots, s_{j,\tau}\}$. On the other hand, a triangulation has its parameter $(\delta_1, \dots, \delta_n)$ and each character δ_i has a τ -Hodge–Tate weight $\text{HT}_\tau(\delta_j)$ (see Section 1.5). The weights match up in that, through (2), we have $\text{HT}_\tau(\delta_j) = s_{j,\tau}$ for all τ and $1 \leq j \leq n$.

Definition 2.2. *Let D be a crystalline (φ, Γ_K) -module such that the τ -Hodge–Tate weights $\{h_{i,\tau}\}$ are distinct for each τ and let P_\bullet be a triangulation of D with parameter $(\delta_1, \dots, \delta_n)$. The critical type of P_\bullet is the collection of permutations $(\sigma_\tau)_\tau$ such that $\text{HT}_\tau(\delta_j) = h_{\sigma_\tau(j),\tau}$ for $j = 1, \dots, n$. We say P_\bullet is non-critical if $\sigma_\tau = \text{id}$ for each τ .*

Remark 2.3. It may be advantageous to see the critical type of a triangulation as lying in the Weyl group of $\text{Res}_{K/\mathbf{Q}_p} \text{GL}_n$ (see [14, 16]).

Example 2.4. Let $(\delta_1, \dots, \delta_n)$ be an n -tuple of continuous characters $\delta_j : \mathbf{Q}_p^\times \rightarrow L^\times$ such that $\text{HT}(\delta_1) < \dots < \text{HT}(\delta_n)$. If $D = \bigoplus_{j=1}^n \mathcal{R}_{\mathbf{Q}_p,L}(\delta_j)$ and $\sigma \in S_n$ then the triangulation

$$(0) \subsetneq \mathcal{R}_{K,L}(\delta_{\sigma(1)}) \subsetneq \dots \subsetneq \bigoplus_{i=1}^j \mathcal{R}_{K,L}(\delta_{\sigma(i)}) \subsetneq \dots \subsetneq D$$

has critical type σ . In particular, only one triangulation of D is non-critical.

2.2. Deformation theory. Continue to let L/\mathbf{Q}_p be a finite extension contained in the fixed algebraic closure $\overline{\mathbf{Q}_p}$. Denote by $\mathfrak{A}\mathfrak{R}_L$ the category of local Artin L -algebras with residue field L . For example, the ring of dual numbers $L[\varepsilon] = L[u]/(u^2)$ is in $\mathfrak{A}\mathfrak{R}_L$. Every element $A \in \mathfrak{A}\mathfrak{R}_L$ is considered a topological ring with the topology defined by its maximal ideal \mathfrak{m}_A and, by definition, morphisms are continuous ring morphisms. A functor $\mathfrak{X} : \mathfrak{A}\mathfrak{R}_L \rightarrow \underline{\text{Set}}$ is (pro)-representable if there exists a complete local noetherian L -algebra $R_{\mathfrak{X}}$ with residue field L and $\text{Hom}_{\text{cont}}(R_{\mathfrak{X}}, A) = \mathfrak{X}(A)$ for all elements $A \in \mathfrak{A}\mathfrak{R}_L$.

Let D denote a (φ, Γ_K) -module over $\mathcal{R}_{K,L}$. A deformation D_A of D to $A \in \mathfrak{A}\mathfrak{R}_L$ is a (φ, Γ_K) -module over $\mathcal{R}_{K,A}$ together with an isomorphism $\pi : D_A \otimes_A L \simeq D$. An isomorphism between deformations (D_A, π) and (D'_A, π') is a (φ, Γ_K) -equivariant isomorphism $\alpha : D_A \xrightarrow{\simeq} D'_A$ such that $\pi' \circ \alpha = \pi$. Thus, we have a functor

$$\mathfrak{X}_D(A) := \{\text{isomorphism classes of deformations of } D \text{ to } A\},$$

which we call the universal deformation functor of D . We have that $\mathfrak{X}_D(L) = \{D\}$ is a single point, and the universal deformation functor of D admits a well-defined, finite-dimensional, Zariski tangent space $\mathfrak{t}_D := \mathfrak{X}_D(L[\varepsilon])$ (see [35, Sections 18 and 23] and [22, Proposition 3.4]).

The tangent space \mathfrak{t}_D admits a canonical description in terms of Galois cohomology. Namely, a deformation \tilde{D} of D to $L[\varepsilon]$ is naturally an extension $0 \rightarrow D \rightarrow \tilde{D} \rightarrow D \rightarrow 0$ in the category of (φ, Γ_K) -modules over $\mathcal{R}_{K,L}$, where the submodule is $\varepsilon\tilde{D}$ and the quotient is $\tilde{D}/\varepsilon\tilde{D}$. The association of \tilde{D} to its extension class defines a canonical L -linear isomorphism

$$\mathfrak{t}_D \simeq \mathrm{Ext}_{(\varphi, \Gamma_K)}^1(D, D) \simeq H^1(\mathrm{ad} D).$$

See [22, Proposition 3.6(ii)] for a direct construction going from \mathfrak{t}_D to $H^1(\mathrm{ad} D)$.

If $\mathfrak{X}' \subset \mathfrak{X}$ is an inclusion of functors on $\mathfrak{A}\mathfrak{R}_L$ then we recall that there is a notion of \mathfrak{X}' being relatively representable over \mathfrak{X} (see [35, Section 19]). Subfunctors $\mathfrak{X}' \subset \mathfrak{X}_D$ are relatively representable if and only if for every morphism of functors $\mathfrak{X}_D \rightarrow \mathfrak{Y}$ with \mathfrak{Y} representable, the base change $\mathfrak{Y}' := \mathfrak{X}' \times_{\mathfrak{X}_D} \mathfrak{Y}$ is representable. In particular being relatively representable is stable under products over \mathfrak{X}_D .

2.3. Deformations of algebraic characters. If A is an affinoid \mathbf{Q}_p -algebra then let $\mathcal{W}(A) = \mathrm{Hom}_{\mathrm{cont}}((\mathcal{O}_K^\times)^n, A^\times)$. This defines a rigid analytic space over \mathbf{Q}_p (a disjoint union of polydiscs) called the p -adic weight space of $\mathrm{GL}_{n/K}$. Since \mathcal{W} is smooth over \mathbf{Q}_p , if $A \in \mathfrak{A}\mathfrak{R}_L$ then the canonical morphism $\mathcal{W}(A) \rightarrow \mathcal{W}(L)$ is surjective. Here we explicitly describe the preimage of \mathbf{Q}_p -algebraic elements of $\mathcal{W}(L)$.

Let $\underline{h} = (h_\tau)_\tau$ be a collection of integers and $z^{\underline{h}} : K^\times \rightarrow L^\times$ be given by

$$z \mapsto \prod_{\tau} \tau(z)^{h_\tau}.$$

The character $z^{\underline{h}}$ has Hodge–Tate weights $(-h_\tau)_\tau$. Every \mathbf{Q}_p -algebraic character of K^\times is $z^{\underline{h}}$ for some \underline{h} . We abuse notation and write \underline{h} for $z^{\underline{h}}$ as well.⁵

We now restrict to \mathcal{O}_K^\times . We have that $\mathcal{O}_K^\times = \mu \times U$ where μ is the torsion subgroup and $U \subset \mathcal{O}_K^\times$ is a finite free \mathbf{Z}_p -submodule. If $\alpha \in \mathcal{O}_K^\times$ we write $\alpha = \omega(\alpha)\langle\alpha\rangle$ where $\langle\alpha\rangle \in U$ and $\omega(\alpha) \in \mu$. Let \underline{h} be as above and suppose $A \in \mathfrak{A}\mathfrak{R}_L$. Suppose that for each τ we choose $\eta_\tau \in A$ such that $\eta_\tau \equiv h_\tau \pmod{\mathfrak{m}_A}$. Then, we define a character $\underline{\eta} : \mathcal{O}_K^\times \rightarrow A^\times$ by

$$z \mapsto \prod_{\tau} \tau(z)^{\eta_\tau} := \prod_{\tau} \exp((\eta_\tau - h_\tau) \cdot \tau(\log\langle z \rangle)) \cdot \tau(z)^{h_\tau}$$

(the exponential converges in A^\times because $\eta_\tau - h_\tau \in \mathfrak{m}_A$). The preimage of \underline{h} under $\mathcal{W}(A) \rightarrow \mathcal{W}(L)$ is exactly the set of characters $\underline{\eta}$.

Note that \underline{h} is a character of K^\times , but extending $\underline{\eta}$ to K^\times requires a choice. After fixing a uniformizer $\varpi_K \in K^\times$, we denote by $\underline{\eta}_{\varpi_K}$ the character of K^\times which acts as $\underline{\eta}$ on \mathcal{O}_K^\times and sends ϖ_K to 1. This defines a rank one (φ, Γ_K) -module $\mathcal{R}_{K,A}(\underline{\eta}_{\varpi_K})$ over $\mathcal{R}_{K,A}$ whose Hodge–Sen–Tate weights are $(-\eta_\tau)_\tau$. The character $\underline{\eta}_{\varpi_K}$ depends on ϖ_K , but if ϖ'_K is another choice of uniformizer then $\underline{\eta}_{\varpi_K} \cdot \underline{\eta}_{\varpi'_K}^{-1}$ is crystalline.

2.4. Hodge–Tate deformations. Let D be a (φ, Γ_K) -module over $\mathcal{R}_{K,L}$. We assume D is Hodge–Tate with distinct Hodge–Tate weights. Let $A \in \mathfrak{A}\mathfrak{R}_L$ and D_A be a deformation of D to A . For each embedding $\tau : K \hookrightarrow L$, the τ -Sen polynomial $P_{D_A}^{\mathrm{Sen}, \tau}(x) \in A[x]$ has distinct roots modulo \mathfrak{m}_A and so $P_{D_A}^{\mathrm{Sen}, \tau}$ completely factors over A by Hensel’s lemma. Thus for each

⁵Warning: if D is a (φ, Γ_K) -module then $D(\underline{h})$ is *not* necessarily a Tate twist.

τ -Hodge–Tate weight $h_\tau \in \mathbf{Z}$ of D there is a unique τ -Hodge–Sen–Tate weight $\eta_\tau \in A$ of D_A such that $\eta_\tau \equiv h_\tau \pmod{\mathfrak{m}_A}$. We say h_τ is a constant weight if $\eta_\tau = h_\tau$. Then, we define

$$\mathfrak{X}_D^{h_\tau}(A) = \{\text{deformations } D_A \text{ of } D \text{ to } A \text{ such that the weight } h_\tau \text{ is constant}\}.$$

By definition this defines a subfunctor $\mathfrak{X}_D^{h_\tau} \subset \mathfrak{X}_D$. If $\{h_{i,\tau}\}$ is the set of all the Hodge–Tate weights of D then the Hodge–Tate deformation functor is defined to be

$$\mathfrak{X}_D^{\text{HT}} := \bigcap_{i,\tau} \mathfrak{X}_D^{h_{i,\tau}}.$$

A deformation D_A is Hodge–Tate if and only if its Hodge–Sen–Tate weights are constant (integers), if and only if it is Hodge–Tate as a (φ, Γ_K) -module over $\mathcal{R}_{K,A}$.

Proposition 2.5. *The subfunctor $\mathfrak{X}_D^{h_{i,\tau}} \subset \mathfrak{X}_D$ is relatively representable for each i, τ . Thus, $\mathfrak{X}_D^{\text{HT}} \subset \mathfrak{X}_D$ is relatively representable as well.*

Proof. Being relatively representable is closed under intersection, so only the first statement needs proving. Observe that $D_A \in \mathfrak{X}_D^{h_{i,\tau}}(A)$ if and only if the Sen operator acts semi-simply on the generalized eigenspace for the eigenvalue $h_{i,\tau}$ inside $D_{\text{Sen}}(D_A)$ (viewing D_A over $\mathcal{R}_{K,L}$). Since this property is closed under subquotients and direct sums, the relative representability of $\mathfrak{X}_D^{h_{i,\tau}}$ follows from Ramakrishna’s criterion [35, Section 25]. \square

2.5. Crystalline deformations. Throughout this section we denote by D a crystalline (φ, Γ_K) -module over $\mathcal{R}_{K,L}$. The fine Selmer group for D , generalizing the corresponding notion for Galois representations [13], is defined by

$$H_f^1(D) := \ker \left(H^1(D) \rightarrow H^1(\Gamma_K, D[t^{-1}]) \right).$$

The cohomology on the right is the continuous cohomology of the profinite group Γ_K . We refer to [38, Section 3.1] and [7, Section 1] for the facts that follow.

First, by [38, Equation 3-2] the dimension of $H_f^1(D)$ is computed by

$$(3) \quad \dim_L H_f^1(D) = \dim_L H^0(D) + \dim_L D_{\text{dR}}(D)/D_{\text{dR}}^+(D).$$

Second, fine Selmer groups arise naturally as tangent spaces to a deformation problem. Indeed, we define the crystalline deformation functor as

$$\mathfrak{X}_{D,f}(A) = \{D_A \in \mathfrak{X}_D(A) : D_A \text{ is crystalline}\}.$$

Since being crystalline is closed under direct sums and subquotients, Ramakrishna’s criterion [35, Section 25] implies that $\mathfrak{X}_{D,f} \subset \mathfrak{X}_D$ is relatively representable. We write $\mathfrak{t}_{D,f} := \mathfrak{X}_{D,f}(L[\varepsilon])$ for the Zariski tangent space to $\mathfrak{X}_{D,f}$.

Since D is crystalline, so is $\text{ad } D$ and the Galois cohomology $H^1(\text{ad } D)$ contains the subspace $H_f^1(\text{ad } D)$. It is explained in [38, Section 3] that an extension class

$$0 \rightarrow D \rightarrow \tilde{D} \rightarrow D \rightarrow 0$$

in $\text{Ext}_{(\varphi, \Gamma_K)}^1(D, D) \simeq H^1(\text{ad } D)$ lies in $H_f^1(\text{ad } D)$ if and only if \tilde{D} is crystalline. If $\text{Ext}_f^1(D, D)$ denotes the corresponding subspace of crystalline extensions then the inclusion $\mathfrak{t}_{D,f} \subset \mathfrak{t}_D$ induces isomorphisms $\mathfrak{t}_{D,f} \simeq \text{Ext}_f^1(D, D) \simeq H_f^1(\text{ad } D)$.

We record here a property of Selmer groups. If D is a crystalline (φ, Γ_K) -module then we write $H_{/f}^1(D) := H^1(D)/H_f^1(D)$.

Lemma 2.6. *If $0 \rightarrow D_1 \rightarrow D_2 \rightarrow D_3 \rightarrow 0$ is a short exact sequence of crystalline (φ, Γ_K) -modules and $H^2(D_1) = 0$ then the canonical morphisms induce a short exact sequence*

$$0 \rightarrow H_{/f}^1(D_1) \rightarrow H_{/f}^1(D_2) \rightarrow H_{/f}^1(D_3) \rightarrow 0.$$

Proof. Consider the commuting diagram

$$\begin{array}{ccccccccc} H^0(D_3) & \longrightarrow & H_f^1(D_1) & \longrightarrow & H_f^1(D_2) & \longrightarrow & H_f^1(D_3) & \longrightarrow & 0 \\ & & \parallel & & \downarrow & & \downarrow & & \\ H^0(D_3) & \longrightarrow & H^1(D_1) & \longrightarrow & H^1(D_2) & \longrightarrow & H^1(D_3) & \longrightarrow & 0. \end{array}$$

The top row is exact by [7, Corollary 1.4.6]. The bottom row is exact because $H^2(D_1) = (0)$. From the snake lemma we get a short exact sequence

$$(4) \quad 0 \rightarrow \frac{\text{coker}(H^0(D_3) \rightarrow H^1(D_1))}{\text{coker}(H^0(D_3) \rightarrow H_f^1(D_1))} \rightarrow H_{/f}^1(D_2) \rightarrow H_{/f}^1(D_3) \rightarrow 0.$$

Since the first term of (4) equals $H_{/f}^1(D_1)$, we are finished. \square

3. WEAKLY-REFINED DEFORMATIONS

3.1. Deforming crystalline eigenvalues. Let D be a crystalline (φ, Γ_K) -module over $\mathcal{R}_{K,L}$. We assume that $h_{1,\tau} = 0$ is the unique least τ -Hodge–Tate weight for each embedding $\tau : K \rightarrow L$. Following Section 2.4 we let $\mathfrak{X}_D^0 = \bigcap_{\tau} \mathfrak{X}_D^{0,\tau}$ be the relatively representable subfunctor of deformations with constant Hodge–Tate weight zero at each embedding.

Suppose that Φ is a crystalline eigenvalue for D (note that it appears in $D_{\text{crys}}(D) = D_{\text{crys}}^+(D) = D^{\Gamma_K}$). Now set

$$(5) \quad \mathfrak{X}_D^{0,\Phi}(A) = \left\{ \begin{array}{l} D_A \in \mathfrak{X}_D^0(A) \text{ such that } D_{\text{crys}}^+(D_A)^{\varphi^{\Gamma_K} = \Phi_A} \text{ is free of rank one} \\ \text{over } K_0 \otimes_{\mathbf{Q}_p} A \text{ for some } \Phi_A \in A^\times \text{ and } \Phi_A \equiv \Phi \pmod{\mathfrak{m}_A} \end{array} \right\}.$$

This clearly defines a subfunctor of $\mathfrak{X}_D^{0,\Phi} \subset \mathfrak{X}_D$.

Proposition 3.1. *If Φ is a simple crystalline eigenvalue for D then $\mathfrak{X}_D^{0,\Phi} \subset \mathfrak{X}_D$ is relatively representable.*

Kisin [31, Proposition 8.13] and Tan [41, Section 5.2] proved Proposition 3.1 for Galois representations. The rest of this subsection is devoted to a proof in the setting of (φ, Γ_K) -modules, adapting [5, Section 2.3]. After twisting D by an unramified character, we may assume that $\Phi = 1$. To emphasize this we write $\mathfrak{X}_D^{0,1} = \mathfrak{X}_D^{0,\Phi=1}$.

If E is a (φ, Γ_K) -module over $\mathcal{R}_{K,L}$, we set

$$F(E) = \{e \in E^{\Gamma_K} : (\varphi^{\Gamma_K} - 1)^n e = (0) \text{ for some } n \geq 1\} \subset D_{\text{crys}}^+(E).$$

The functor $F(-)$ is left exact, and since $\Phi = 1$ is a simple eigenvalue of D we have $\dim_L F(D) = (K_0 : \mathbf{Q}_p)$.

Let A denote an element in $\mathfrak{A}\mathfrak{R}_L$. If D_A is a (φ, Γ_K) -module over $\mathcal{R}_{K,A}$ and M is an A -module then we consider $D_A \otimes_A M$ as a (φ, Γ_K) -module over $\mathcal{R}_{K,L}$ with the trivial actions on M . The L -vector space $F(D_A \otimes_A M)$ is naturally equipped with the structure of an A -module. We let ℓ_A denote the length function on A -modules.

Lemma 3.2. *Let $A \in \mathfrak{A}\mathfrak{R}_L$ and $D_A \in \mathfrak{X}_D(A)$. If M is a finite length A -module then $\ell_A(F(D_A \otimes_A M)) \leq (K_0 : \mathbf{Q}_p)\ell_A(M)$. In particular, if $I \subset A$ is an ideal then $\ell_A(F(ID_A)) \leq (K_0 : \mathbf{Q}_p)\ell_A(I)$.*

Proof. The second statement follows from the first since D_A is flat over A (it is even free), so $ID_A \simeq D_A \otimes_A I$. The first statement is an immediate dévissage using the left-exactness of $F(-)$ and the fact explained above that $\ell_A(F(D)) = \dim_L F(D) = (K_0 : \mathbf{Q}_p)$. \square

Lemma 3.3. *Let $D_A \in \mathfrak{X}_D(A)$. The following are equivalent*

- (a) $D_A \in \mathfrak{X}_D^{0, \Phi=1}(A)$.
- (b) $F(D_A)$ is free of rank one over $K_0 \otimes_{\mathbf{Q}_p} A$.
- (c) $\ell_A(F(D_A)) = (K_0 : \mathbf{Q}_p)\ell_A(A)$.
- (d) The natural map $F(D_A)/\mathfrak{m}_A F(D_A) \rightarrow F(D)$ is an isomorphism.

Proof. First assume that $D_A \in \mathfrak{X}_D^{0, \Phi=1}(A)$ and let $\Phi_A \in A$ be the deformation of $\Phi = 1$ as in the definition (5). Then $D_{\text{crys}}^+(D_A)^{\varphi^{fK}=\Phi_A}$ is a free, rank one, $K_0 \otimes_{\mathbf{Q}_p} A$ -submodule of $F(D_A)$. It cannot be proper since that would imply $\ell_A(F(D_A)) > \ell_A(D_{\text{crys}}^+(D_A)^{\varphi^{fK}=\Phi_A}) = \ell_A(K_0 \otimes_{\mathbf{Q}_p} A)$, which would contradict Lemma 3.2 (applied with $M = A$). Thus, $F(D_A) = D_{\text{crys}}^+(D_A)^{\varphi^{fK}=\Phi_A}$ is free of rank one over $K_0 \otimes_{\mathbf{Q}_p} A$. This proves (a) implies (b).

We clearly have (b) implies (c). Now we show (c) implies (d). Since $F(-)$ is left exact we have an exact sequence

$$(6) \quad 0 \rightarrow F(\mathfrak{m}_A D_A) \rightarrow F(D_A) \rightarrow F(D).$$

Under the assumption (c), considering the lengths in (6), Lemma 3.2 implies that $F(D_A) \rightarrow F(D)$ is surjective. But then it follows that $F(D_A)/\mathfrak{m}_A F(D_A) \rightarrow F(D)$ is onto as well. Since the two A -modules have the same length, we've proven (d).

It remains to prove (d) implies (a) (we will simultaneously show (d) implies (b)). First, we can choose a vector $v \in F(D_A)$ such that the image in $F(D)$ is a $K_0 \otimes_{\mathbf{Q}_p} L$ -module basis. It follows that the $K_0 \otimes_{\mathbf{Q}_p} A$ -module spanned by v inside $F(D_A)$ is free of rank one.⁶ By considering lengths, Lemma 3.2 implies that the containment $(K_0 \otimes_{\mathbf{Q}_p} A) \cdot v \subset F(D_A)$ is necessarily an equality. This shows (d) implies (b) and since $F(D_A)$ is φ -stable, $\varphi(v) = xv$ for some $x \in K_0 \otimes_{\mathbf{Q}_p} A$. But then, $\varphi^{fK}(v) = x\varphi(x) \cdots \varphi^{fK-1}(x)v$ where the φ on the right-hand side is the natural A -linear Frobenius on $K_0 \otimes_{\mathbf{Q}_p} A$. Since $x\varphi(x) \cdots \varphi^{fK-1}(x)$ lies in $A^\times \subset K_0 \otimes_{\mathbf{Q}_p} A$ we will call it Φ_A . Note that since v spans $F(D)$ modulo \mathfrak{m}_A , we know that $\Phi_A \equiv 1 \pmod{\mathfrak{m}_A}$. Finally, since $v \in F(D_A) \subset D_A^{\Gamma_K} = D_{\text{crys}}^+(D_A)$ as well, we've shown that $v \in D_{\text{crys}}^+(D_A)^{\varphi^{fK}=\Phi_A}$. But then we have $(K_0 \otimes_{\mathbf{Q}_p} A) \cdot v = D_{\text{crys}}^+(D_A)^{\varphi^{fK}=\Phi_A} = F(D_A)$ and we've shown (d) implies (a). \square

We are now ready to prove Proposition 3.1. If $D_A \in \mathfrak{X}_D(A)$ and $A \rightarrow A'$ is an arrow in $\mathfrak{A}\mathfrak{R}_L$ then we write $D_{A'} := D_A \otimes_A A'$.

Proof of Proposition 3.1 with $\Phi = 1$. To show $\mathfrak{X}_D^{0, \Phi=1} \subset \mathfrak{X}_D$ is relatively representable we will use Schlessinger's criterion [35, Section 23], labeled by conditions (1), (2), and (3) in *loc.*

⁶This is a minor modification of an argument given by Bellaïche and Chenevier in the proof of [5, Proposition 2.3.9]. To reduce to their argument, use that the image of v is non-zero in $F(D) \otimes_{K_0, \tau} L$ for each embedding $\tau : K_0 \hookrightarrow L$.

cit. Condition (1) is clear. Given (1) and (3) in *loc. cit.* it is enough to check condition (2) with $C = L$, but that it is also clear (compare with [31, Propositions 8.13 and 8.7]).

It remains to check condition (3): if $A \subset A'$ is an inclusion in $\mathfrak{A}\mathfrak{R}_L$, $D_A \in \mathfrak{X}_D(A)$ and $D_{A'} \in \mathfrak{X}_D^{0, \Phi=1}(A')$ then $D_A \in \mathfrak{X}_D^{0, \Phi=1}(A)$. We start with the assumptions of condition (3) and consider the exact sequence $0 \rightarrow F(D_A) \rightarrow F(D_{A'}) \rightarrow F(D_A \otimes_A A'/A)$. Computing lengths,

$$(7) \quad \begin{aligned} \ell_A(F(D_{A'})) - (K_0 : \mathbf{Q}_p)\ell_A(A'/A) &\leq \ell_A(F(D_{A'})) - \ell_A(F(D_A \otimes_A A'/A)) \\ &\leq \ell_A(F(D_A)) \end{aligned}$$

Here we used Lemma 3.2 for the first inequality. Since $D_{A'} \in \mathfrak{X}_D^{0, \Phi=1}(A')$ we have that $F(D_{A'})$ is free of rank one over $K_0 \otimes_{\mathbf{Q}_p} A'$ and thus (7) implies $(K_0 : \mathbf{Q}_p)\ell_A(A) \leq \ell_A(F(D_A))$. The reverse inequality is also true by Lemma 3.2. So, we have shown that $\ell_A(F(D_A)) = (K_0 : \mathbf{Q}_p)\ell_A(A) = \ell_A(K_0 \otimes_{\mathbf{Q}_p} A)$. But then $D_A \in \mathfrak{X}_D^{0, \Phi=1}(A)$ by Lemma 3.3. \square

3.2. Weakly-refined deformations. We temporarily fix a uniformizer $\varpi_K \in K^\times$. We also assume D is a crystalline (φ, Γ_K) -module over $\mathcal{R}_{K,L}$ and that for each embedding $\tau : K \hookrightarrow L$ there is a unique least τ -Hodge–Tate weight $h_{1,\tau}$.

If $A \in \mathfrak{A}\mathfrak{R}_L$ and D_A is a deformation of D to A we write $\eta_{1,\tau}$ for the Hodge–Sen–Tate weight of D_A satisfying $\eta_{1,\tau} \equiv h_{1,\tau} \pmod{\mathfrak{m}_A}$ (cf. Section 2.4). Then, by Section 2.3 we can construct a character $\underline{\eta}_{1,\varpi_K} : K^\times \rightarrow A^\times$ whose value on ϖ_K is 1 and whose τ -Hodge–Sen–Tate weight is $\eta_{1,\tau}$ for each embedding $\tau : K \hookrightarrow L$. The twisting operation $D_A \mapsto D_A(\underline{\eta}_{1,\varpi_K})$ defines a natural transformation $\mathfrak{X}_D \rightarrow \mathfrak{X}_{D(h_{1,\varpi_K})}^0$. (Fixing ϖ_K makes the twisting functorial.)

Continue with the assumptions of the previous paragraphs. If ϕ is a simple crystalline eigenvalue for D then $\Phi := \phi \prod_\tau \tau(\varpi_K)^{-h_{1,\tau}}$ is a simple eigenvalue for $D(h_{1,\varpi_K})$. Note that the least Hodge–Tate weight of $D(h_{1,\varpi_K})$ is zero at each embedding τ . We now define a functor \mathfrak{X}_D^ϕ as a fibered product

$$\begin{array}{ccc} \mathfrak{X}_D^\phi & \longrightarrow & \mathfrak{X}_D \\ \downarrow & & \downarrow D_A \mapsto D_A(\underline{\eta}_{1,\varpi_K}) \\ \mathfrak{X}_{D(h_{1,\varpi_K})}^{0,\Phi} & \longrightarrow & \mathfrak{X}_{D(h_{1,\varpi_K})}^0 \end{array}$$

where the bottom arrow is the natural inclusion.

Since being relatively representable is stable under base change, Proposition 3.1 implies that $\mathfrak{X}_D^\phi \subset \mathfrak{X}_D$ is relatively representable. Furthermore, \mathfrak{X}_D^ϕ is independent of ϖ_K : the twisting $D_A \mapsto D_A(\underline{\eta}_{1,\varpi_K})$ and Φ depend on the choice but the dependencies cancel.⁷

For the rest of this subsection, we assume that D is a crystalline (φ, Γ_K) -module over $\mathcal{R}_{K,L}$ with distinct Hodge–Tate weights and distinct crystalline eigenvalues. We also equip D with a triangulation P_\bullet whose parameter we write $(\delta_1, \dots, \delta_n)$. We denote by (ϕ_1, \dots, ϕ_n) the corresponding list of crystalline eigenvalues given by (2).

If $1 \leq j \leq n$ then consider the j th exterior power $\wedge^j D$ of D . If D_A is a deformation of D to A then $\wedge^j D_A$ is a (φ, Γ_K) -module over $\mathcal{R}_{K,A}$ which deforms $\wedge^j D$. If the τ -Hodge–Tate

⁷As we mentioned at the end of Section 2.3, if ϖ_K and ϖ'_K are two different uniformizers then $\underline{\eta}_{1,\varpi_K}$ and $\underline{\eta}_{1,\varpi'_K}$ differ by a crystalline character. Moreover, the crystalline eigenvalue exactly matches the difference between Φ and the corresponding Φ' .

weights of D are $h_{1,\tau} < h_{2,\tau} < \dots < h_{n,\tau}$ then the τ -Hodge–Tate weight $h_{1,\tau} + \dots + h_{j,\tau}$ for $\wedge^j D$ is the unique least τ -Hodge–Tate weight for each τ . Moreover, if $1 \leq j \leq n$ then $\phi_1 \cdots \phi_j$ is an eigenvalue for φ^{f_K} acting on $D_{\text{crys}}(\wedge^j D)$.

Definition 3.4. A triangulation P_\bullet is called *regular* if $\phi_1 \cdots \phi_j$ is a simple crystalline eigenvalue in $D_{\text{crys}}(\wedge^j D)$ for $1 \leq j \leq n$. We say P_\bullet is *regular generic* if P_\bullet is regular and in addition $H^2(\delta_i \delta_j^{-1}) = (0)$ for all $1 \leq i, j \leq n$.

Remark 3.5. If P_\bullet is a regular triangulation as in Definition 3.4 then the crystalline eigenvalues of D are necessarily distinct. Since $H^0(\delta_i \delta_j^{-1}) \subset D_{\text{crys}}(\mathcal{R}_{K,L}(\delta_i \delta_j^{-1}))^{\varphi=1}$, it follows that $H^0(\delta_i \delta_j^{-1}) = (0)$ when $i \neq j$.

Definition 3.6. Let D be a crystalline (φ, Γ_K) -module with distinct Hodge–Tate weights. If P_\bullet is a regular triangulation then the weakly-refined deformation functor with respect to P_\bullet is the fibered product $\mathfrak{X}_{D,P_\bullet}^{\text{Ref}}$ given by

$$\begin{array}{ccc} \mathfrak{X}_{D,P_\bullet}^{\text{Ref}} & \longrightarrow & \mathfrak{X}_D \\ \downarrow & & \downarrow^{D_A \mapsto (\wedge^j D_A)_j} \\ \prod_{j=1}^n \mathfrak{X}_{\wedge^j D}^{\phi_1 \cdots \phi_j} & \longrightarrow & \prod_{j=1}^n \mathfrak{X}_{\wedge^j D} \end{array}$$

where the bottom horizontal arrow is the natural inclusion.

When P_\bullet is a regular triangulation, Proposition 3.1 implies that each arrow $\mathfrak{X}_{\wedge^j D}^{\phi_1 \cdots \phi_j} \rightarrow \mathfrak{X}_{\wedge^j D}$ is relatively representable. Thus, so is $\mathfrak{X}_{D,P_\bullet}^{\text{Ref}} \subset \mathfrak{X}_D$.

3.3. A constant weight tangent space. Our goal for the rest of Section 3 is to compute the Zariski tangent spaces $\mathfrak{t}_{D,P_\bullet}^{\text{Ref}}$ to functors of the form $\mathfrak{X}_{D,P_\bullet}^{\text{Ref}}$. This particular subsection concerns the constant weight subfunctor $\mathfrak{X}_{D,P_\bullet}^{\text{Ref,HT}} := \mathfrak{X}_{D,P_\bullet}^{\text{Ref}} \cap \mathfrak{X}_D^{\text{HT}}$, which admits a simpler description: if D_A is a Hodge–Tate deformation of D then the twisting character(s) denoted $\frac{\eta_{1,\varphi_K}}{(8)}$ are crystalline, and thus the points of $\mathfrak{X}_D^{\text{Ref,HT}}$ are given by

$$\mathfrak{X}_D^{\text{Ref,HT}}(A) = \left\{ \begin{array}{l} D_A \in \mathfrak{X}_D^{\text{HT}}(A) \text{ such that for some collection } \phi_{j,A} \equiv \phi_j \pmod{\mathfrak{m}_A}, \\ D_{\text{crys}}(\wedge^j D_A)^{\varphi^{f_K} = \phi_{1,A} \cdots \phi_{j,A}} \text{ is free of rank one over } K_0 \otimes_{\mathbb{Q}_p} A \text{ for} \\ \text{each } 1 \leq j \leq n. \end{array} \right\}.$$

From the description (8) it is clear that $\mathfrak{X}_{D,P_\bullet}^{\text{Ref,HT}}$ contains the crystalline deformation functor $\mathfrak{X}_{D,f}$ for any P_\bullet . In Theorem 3.12 below we compute a bound for the dimension of quotient spaces of the form $\mathfrak{t}_{D,P_\bullet}^{\text{Ref,HT}} / \mathfrak{t}_{D,f}$. Let us preview our theorem by remarking on the non-critical case.

Remark 3.7. Suppose that D is a crystalline (φ, Γ_K) -module equipped with a regular generic triangulation P_\bullet which is *non-critical*. Then, every weakly-refined deformation is trianguline [5, Theorem 2.5.6] and every trianguline deformation of constant Hodge–Tate weight is crystalline [5, Theorem 2.5.1]. Thus $\mathfrak{t}_{D,P_\bullet}^{\text{Ref,HT}} / \mathfrak{t}_{D,f} = (0)$ in this case.

The next three lemmas expand on the previous remark for triangulations that are possibly critical, and our proof of Theorem 3.12 gives a new proof of the remark.

Lemma 3.8. *Suppose that D is a (φ, Γ_K) -module over $\mathcal{R}_{K,L}$ and $\iota : P_1 \hookrightarrow D$ is a saturated (φ, Γ_K) -submodule. Let $Q_1 := \text{coker}(\iota)$. Consider the canonical morphisms*

$$\begin{array}{ccc} \text{Ext}_{(\varphi, \Gamma_K)}^1(Q_1, D) & \xrightarrow{f} & \text{Ext}_{(\varphi, \Gamma_K)}^1(D, D) \\ & & \downarrow h \\ \text{Ext}_{(\varphi, \Gamma_K)}^1(Q_1, Q_1) & & \end{array}$$

- (a) *If $\tilde{D} = f(X)$ for some $X \in \text{Ext}_{(\varphi, \Gamma_K)}^1(Q_1, D)$ then there exists a saturated embedding $P_1 \otimes_{\mathcal{R}_{K,L}} \mathcal{R}_{K,L[\varepsilon]} \xrightarrow{\tilde{\iota}} \tilde{D}$ such that $\tilde{\iota} \equiv \iota \pmod{\varepsilon}$.*
- (b) *If $\dim_L \text{Hom}_{(\varphi, \Gamma_K)}(P_1, D) = 1$ and $\tilde{D} = f(X)$ for some $X \in \text{Ext}_{(\varphi, \Gamma_K)}^1(Q_1, D)$ then $\tilde{\iota}$ in part (a) is unique up to a scalar $1 + \mu\varepsilon \in L[\varepsilon]^\times$ and $h(X) = \text{coker}(\tilde{\iota})$.*

Proof. We first recall the definitions of f and h . Let

$$0 \rightarrow D \xrightarrow{j_X} X \xrightarrow{\pi_X} Q_1 \rightarrow 0$$

be an extension in $\text{Ext}_{(\varphi, \Gamma_K)}^1(Q_1, D)$. Then, the embedding $\iota : P_1 \hookrightarrow D$ induces an embedding $j_X \circ \iota : P_1 \hookrightarrow X$ and $h(X) := \text{coker}(j_X \circ \iota)$. If $\theta : D \rightarrow Q_1$ is the quotient map then $f(X)$ is defined by

$$f(X) := \ker \left(X \oplus D \xrightarrow{(\pi_X, -\theta)} Q_1 \right) = \{(x, d) \in X \oplus D : \pi_X(x) = \theta(d)\}.$$

The (φ, Γ_K) -module structure on $f(X)$ is coordinate-wise, and the structure of multiplication by ε on $f(X)$ is given by $\varepsilon \cdot (x, d) = (j_X(d), 0)$.

Now let's prove (a). Let $X \in \text{Ext}_{(\varphi, \Gamma_K)}^1(Q_1, D)$ and $\tilde{D} = f(X)$. We have a canonical (φ, Γ_K) -equivariant embedding $P_1 \hookrightarrow D \hookrightarrow X \oplus D$ given by $\tilde{\iota}_0(p) = (0, \iota(p))$. Since $\pi_X(0) = 0 = \theta(\iota(p))$ we see that $\tilde{\iota}_0(P_1) \subset \tilde{D}$. We define $\tilde{\iota}$ on $P_1 \otimes_{\mathcal{R}_{K,L}} \mathcal{R}_{K,L[\varepsilon]}$ by $\tilde{\iota}(p) = \tilde{\iota}_0(p)$ and $\tilde{\iota}(\varepsilon p) = (j_X(\iota(p)), 0)$ for each $p \in P_1$. Note that this is well-defined because

$$\varepsilon \tilde{\iota}(p) = \varepsilon \cdot (0, \iota(p)) = (j_X(\iota(p)), 0) = \tilde{\iota}(\varepsilon p).$$

Moreover, $\tilde{\iota}$ is injective because $\text{im}(\tilde{\iota}_0) \cap \varepsilon \tilde{D} = (0)$. It is clear that $\tilde{\iota} \pmod{\varepsilon} = \iota$ and $\text{im}(\tilde{\iota})$ is saturated by Lemma 2.1.

We now prove part (b). A general $\tilde{\iota}$ is determined by $\tilde{\iota}(p)$ with $p \in P_1$ by $L[\varepsilon]$ -linearity. Since $\tilde{\iota} \equiv \iota \pmod{\varepsilon}$ we may write $\tilde{\iota}(p) = (j_X(\beta(p)), \iota(p))$ where $\beta : P_1 \rightarrow D$ is a (φ, Γ_K) -equivariant map. By assumption, β is of the form $\mu \cdot \iota$ for some $\mu \in L$ and so $\tilde{\iota}(p) = (1 + \mu\varepsilon) \cdot \tilde{\iota}_0(p)$. This shows that $\tilde{\iota}$ is unique up to a scalar $1 + \mu\varepsilon \in L[\varepsilon]^\times$.

It remain to show $h(X) = \text{coker}(\tilde{\iota})$. By the previous paragraph, we may assume $\tilde{\iota}$ is explicitly given as in the proof of (a). First note the map $\tilde{D} = f(X) \rightarrow X$ given by $(x, d) \mapsto x$ is surjective. Furthermore, if $p, q \in P_1$ then $\tilde{\iota}(p + q\varepsilon) = (j_X \iota(q), \iota(p)) \mapsto j_X \iota(q) \in (j_X \circ \iota)(P_1)$. Thus we have a well-defined (φ, Γ_K) -equivariant surjection $\alpha : \tilde{D} / \text{im}(\tilde{\iota}) \twoheadrightarrow X / (j_X \circ \iota)(P_1)$. Suppose that $\tilde{d} \in \ker(\alpha)$ and write $\tilde{d} = (j_X(\iota(p)), d)$ for some $p \in P_1$ and $d \in D$. Since $\theta(d) = \pi_X(j_X(\iota(p))) = 0$ we may write $d = \iota(p')$ for some $p' \in P_1$. But then $\tilde{d} = \tilde{\iota}(p' + \varepsilon p) \in \text{im}(\tilde{\iota})$. This shows α is injective and so $h(X) = X / (j_X \circ \iota)(P_1) = \text{coker}(\tilde{\iota})$. \square

Lemma 3.9. *Let D be a crystalline (φ, Γ_K) -module over $\mathcal{R}_{K,L}$ and let $\mathcal{R}_{K,L}(\delta_1) \subset D$ be a saturated rank one (φ, Γ_K) -submodule. Let ϕ_1 be the crystalline eigenvalue appearing in*

$D_{\text{crys}}(\mathcal{R}_{K,L}(\delta_1))$, assume that ϕ_1 is simple in $D_{\text{crys}}(D)$ and let $h_1 = (h_{1,\tau})$ be the Hodge–Tate weights of δ_1 . Let $\mathfrak{t}_D^{\phi_1, h_1}$ be the Zariski tangent space to $\mathfrak{X}_D^{\phi_1, h_1} := \mathfrak{X}_D^{\phi_1} \cap \bigcap_{\tau} \mathfrak{X}_D^{h_{1,\tau}}$ as in Sections 2.4 and 3.2.

- (a) The composition $\mathfrak{t}_D^{\phi_1, h_1} \hookrightarrow H^1(\text{ad } D) \rightarrow H_{/f}^1(\text{ad } D) \rightarrow H_{/f}^1(D(\delta_1^{-1}))$ is zero.
- (b) If $Q_1 = \text{coker}(\mathcal{R}_{K,L}(\delta_1) \rightarrow D)$ and $H^2(D \otimes Q_1^\vee) = (0)$ then there is a natural embedding $\mathfrak{t}_D^{\phi_1, h_1} / \mathfrak{t}_{D,f} \hookrightarrow H_{/f}^1(D \otimes Q_1^\vee)$.

Proof. Suppose that $\tilde{D} \in \mathfrak{t}_D^{\phi_1, h_1}$. The least Hodge–Tate weight $h_{1,\tau}$ is constant in \tilde{D} for each embedding $\tau : K \hookrightarrow L$. Thus there exists a $\tilde{\phi}_1 \in L[\varepsilon]^\times$ deforming ϕ_1 such that $D_{\text{crys}}(\tilde{D})^{\varphi^{fK} = \tilde{\phi}_1}$ is free of rank one over $K_0 \otimes_{\mathbf{Q}_p} L[\varepsilon]$ (compare with (8)). By [9, Lemma 7.2], the image of \tilde{D} under the natural map $\text{Ext}_{(\varphi, \Gamma_K)}^1(D, D) \rightarrow \text{Ext}_{(\varphi, \Gamma_K)}^1(\mathcal{R}_{K,L}(\delta_1), D)$ lands inside the subspace $\text{Ext}_f^1(\mathcal{R}_{K,L}(\delta_1), D)$ of crystalline extensions (the reference is valid because we have assumed that ϕ_1 is a simple eigenvalue in $D_{\text{crys}}(D)$). This proves part (a).

To prove part (b), we apply Lemma 2.6 to the short exact sequence

$$(9) \quad 0 \rightarrow D \otimes Q_1^\vee \rightarrow \text{ad } D \rightarrow D(\delta_1^{-1}) \rightarrow 0$$

of (φ, Γ_K) -modules. The hypotheses of Lemma 2.6 are satisfied because D , and thus each term in (9), is crystalline and $H^2(D \otimes Q_1^\vee) = (0)$ by assumption in this lemma. We conclude from Lemma 2.6 that there is a natural short exact sequence

$$(10) \quad 0 \rightarrow H_{/f}^1(D \otimes Q_1^\vee) \rightarrow H_{/f}^1(\text{ad } D) \rightarrow H_{/f}^1(D(\delta_1^{-1})) \rightarrow 0.$$

The subspace $\mathfrak{t}_D^{\phi_1, h_1} / \mathfrak{t}_{D,f} \subset H_{/f}^1(\text{ad } D)$ maps to zero in the final term of (10) by part (a) of this lemma. This proves (b). \square

Lemma 3.10. *Let D be a crystalline (φ, Γ_K) -module equipped with a regular generic triangulation P_\bullet and let $\mathcal{R}_{K,L}(\delta_1) = P_1$. Set $Q_1 = \text{coker}(\mathcal{R}_{K,L}(\delta_1) \rightarrow D)$ and let P'_\bullet be the induced triangulation on Q_1 . The composition*

$$\mathfrak{t}_{D, P_\bullet}^{\text{Ref, HT}} / \mathfrak{t}_{D,f} \hookrightarrow H_{/f}^1(D \otimes Q_1^\vee) \rightarrow H_{/f}^1(\text{ad } Q_1)$$

has image inside the subspace $\mathfrak{t}_{Q_1, P'_\bullet}^{\text{Ref, HT}} / \mathfrak{t}_{Q_1,f} \subset H_{/f}^1(\text{ad } Q_1)$.

Remark 3.11. There are minor clarifications needed for the lemma. First, the composition in Lemma 3.10 is well-defined by Lemma 3.9(b) and the inclusion $\mathfrak{t}_{D, P_\bullet}^{\text{Ref, HT}} \subset \mathfrak{t}_D^{\phi_1, h_1}$ (the notation as in Lemma 3.9). Second, the induced triangulation P'_\bullet on Q_1 is regular generic because P_\bullet itself is regular generic. This gives content to the conclusion of Lemma 3.10.

Proof of Lemma 3.10. First note that $H^2(\delta_j \delta_i^{-1}) = (0)$ for each i, j because P_\bullet is assumed to be regular generic. In particular, by the long exact sequence in cohomology we deduce that $H^2(D \otimes Q_1^\vee) = (0)$.

Write α for the composition $\mathfrak{t}_{D, P_\bullet}^{\text{Ref, HT}} / \mathfrak{t}_{D,f} \rightarrow H_{/f}^1(D \otimes Q_1^\vee) \rightarrow H_{/f}^1(\text{ad } Q_1)$. Now suppose that $[\tilde{D}] \in \mathfrak{t}_{D, P_\bullet}^{\text{Ref, HT}} / \mathfrak{t}_{D,f}$. The tangent space $\mathfrak{t}_{D, P_\bullet}^{\text{Ref, HT}}$ is contained in the tangent space $\mathfrak{t}_D^{\phi_1, h_1}$ as in Lemma 3.9. Since $H^2(D \otimes Q_1^\vee) = (0)$, we may use Lemma 3.9(b) to choose a representative $\tilde{D} \in \mathfrak{t}_{D, P_\bullet}^{\text{Ref, HT}}$ such that \tilde{D} is in the image of $H^1(D \otimes Q_1^\vee) \rightarrow H^1(\text{ad } D)$.

By Lemma 3.8(a) there exists a constant deformation $\tilde{\tau} : \mathcal{R}_{K,L[\varepsilon]}(\delta_1) \hookrightarrow \tilde{D}$ which is a direct summand over $\mathcal{R}_{K,L[\varepsilon]}$. The space $H^0(D(\delta_1^{-1}))$ is one-dimensional since P_\bullet is regular (see the

remark following Definition 3.4) and so Lemma 3.8(b) implies that

$$\alpha([\tilde{D}]) = \text{coker}(\tilde{\iota}) \bmod H_f^1(\text{ad } Q_1).$$

If $\tilde{Q}_1 := \text{coker}(\tilde{\iota}) \in H^1(\text{ad } Q_1)$ then \tilde{Q}_1 has constant Hodge–Tate weights because \tilde{D} has constant Hodge–Tate weights. It suffices now to show that \tilde{Q}_1 is a weakly-refined deformation with respect to the triangulation P'_\bullet .

Let $\tilde{\phi}_1, \dots, \tilde{\phi}_n$ be the elements of $L[\varepsilon]^\times$ witnessing \tilde{D} as being weakly-refined (see (8)). Note $\tilde{\phi}_1$ must be equal to ϕ_1 because ϕ_1 is a simple eigenvalue in $D_{\text{crys}}(D)$ and $\tilde{\iota}$ witnesses ϕ_1 appearing in $D_{\text{crys}}(\tilde{D})$. Now consider $2 \leq m \leq n$ and the short exact sequence of (φ, Γ_K) -modules over $\mathcal{R}_{K,L[\varepsilon]}$

$$0 \rightarrow \wedge^{m-1}(\tilde{Q}_1) \otimes_{\mathcal{R}_{K,L[\varepsilon]}} \mathcal{R}_{K,L[\varepsilon]}(\delta_1) \rightarrow \wedge^m \tilde{D} \rightarrow \wedge^m \tilde{Q}_1 \rightarrow 0.$$

Recall that $D_{\text{crys}}(-)$ is left exact. Since the eigenvalue $\phi_1 \dots \phi_m$ appearing in $D_{\text{crys}}(\wedge^m D)$ does not appear in $D_{\text{crys}}(\wedge^m Q_1)$, by the regular condition on P_\bullet , we conclude that for $m = 2, \dots, n$ there is a natural equality

$$D_{\text{crys}} \left(\wedge^{m-1}(\tilde{Q}_1) \otimes_{\mathcal{R}_{K,L[\varepsilon]}} \mathcal{R}_{K,L[\varepsilon]}(\delta_1) \right)^{\varphi^{fK} = \phi_1 \tilde{\phi}_2 \dots \tilde{\phi}_m} \xrightarrow{\cong} D_{\text{crys}} \left(\wedge^m(\tilde{D}) \right)^{\varphi^{fK} = \phi_1 \tilde{\phi}_2 \dots \tilde{\phi}_m}.$$

Twisting the left hand side by the constant crystalline (φ, Γ_K) -module $\mathcal{R}_{K,L[\varepsilon]}(\delta_1^{-1})$ we have shown that $\tilde{Q}_1 \in \mathfrak{t}_{Q_1, P'_\bullet}^{\text{Ref}}$. \square

We now give an upper bound for the constant weight, weakly-refined, deformations, up to the crystalline deformations. We restate our hypotheses for clarity.

Theorem 3.12. *If D is a crystalline (φ, Γ_K) -module over $\mathcal{R}_{K,L}$ equipped with a regular generic triangulation P_\bullet with parameter $(\delta_1, \dots, \delta_n)$ then*

$$(11) \quad \dim_L \mathfrak{t}_{D, P_\bullet}^{\text{Ref, HT}} / \mathfrak{t}_{D, f} \leq \sum_{1 \leq i < j \leq n} \dim_L H_{/f}^1(\delta_i \delta_j^{-1}).$$

Proof. We argue by induction on n . If $n = 1$ then $\mathfrak{t}_{D, P_\bullet}^{\text{Ref}} = \mathfrak{t}_D$ and $\mathfrak{t}_{D, P_\bullet}^{\text{Ref, HT}} = \mathfrak{t}_{D, f}$. Thus the bound is true (and an equality) in this case.

Now suppose that $n > 1$ and let $Q_1 = \text{coker}(\mathcal{R}_{K,L}(\delta_1) \rightarrow D)$. Then $H^2(Q_1^\vee(\delta_1)) = (0)$ because P_\bullet is regular generic. By Lemmas 2.6 and 3.9 we have a diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & U & \longrightarrow & \mathfrak{t}_{D, P_\bullet}^{\text{Ref, HT}} / \mathfrak{t}_{D, f} & \longrightarrow & V \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H_{/f}^1(Q_1^\vee(\delta_1)) & \longrightarrow & H_{/f}^1(D \otimes Q_1^\vee) & \longrightarrow & H_{/f}^1(\text{ad } Q_1) \longrightarrow 0 \end{array}$$

where U and V are defined by the diagram itself. We separately bound $\dim_L U$ and $\dim_L V$.

By Lemma 3.10, $V \subset \mathfrak{t}_{Q_1, P'_\bullet}^{\text{Ref, HT}} / \mathfrak{t}_{Q_1, f}$ where P'_\bullet is the triangulation induced on Q_1 from P_\bullet . Thus by induction we have

$$\dim_L V \leq \dim_L \mathfrak{t}_{Q_1, P'_\bullet}^{\text{Ref, HT}} / \mathfrak{t}_{Q_1, f} \leq \sum_{2 \leq i < j \leq n} \dim_L H_{/f}^1(\delta_i \delta_j^{-1}).$$

On the other hand, $U \subset H_{/f}^1(Q_1^\vee(\delta_1))$ and so

$$\dim_L U \leq \dim_L H_{/f}^1(Q_1^\vee(\delta_1)) \stackrel{\text{Lemma 2.6}}{=} \sum_{j=2}^n \dim_L H_{/f}^1(\delta_1 \delta_j^{-1}).$$

Putting the two upper bounds together we get the result. \square

Finally we translate Theorem 3.12 into an upper bound in terms of critical types.

Lemma 3.13. *If $\delta : K^\times \rightarrow L^\times$ is a crystalline character and $H^2(\delta) = (0)$ then $\dim_L H_{/f}^1(\delta) = \#\{\tau : K \hookrightarrow L : \text{HT}_\tau(\delta) \geq 0\}$.*

Proof. By the formula (3) we have that

$$\dim_L H_{/f}^1(\delta) = \dim_L H^0(\delta) + \dim_L D_{\text{dR}}(\mathcal{R}_{K,L}(\delta)) / D_{\text{dR}}^+(\mathcal{R}_{K,L}(\delta)).$$

On the other hand, since $H^2(\delta) = (0)$, the Euler–Poincaré characteristic formula [33, Theorem 1.2(a)] implies that $\dim_L H^1(\delta) = (K : \mathbf{Q}_p) + \dim_L H^0(\delta)$. Thus,

$$\begin{aligned} \dim_L H_{/f}^1(\delta) &= (K : \mathbf{Q}_p) - \dim_L D_{\text{dR}}(\mathcal{R}_{K,L}(\delta)) / D_{\text{dR}}^+(\mathcal{R}_{K,L}(\delta)) \\ &= (K : \mathbf{Q}_p) - \#\{\tau : \text{HT}_\tau(\delta) < 0\}. \end{aligned}$$

The result is now clear. \square

Recall that if $\sigma \in S_n$ then its length is given by $\ell(\sigma) = \#\{(i, j) : i < j \text{ and } \sigma(i) > \sigma(j)\}$.

Corollary 3.14. *Suppose that D is a crystalline (φ, Γ_K) -module over $\mathcal{R}_{K,L}$ equipped with a regular generic triangulation P_\bullet and let $(\sigma_\tau)_\tau$ be the critical type of P_\bullet . Then*

$$\dim_L \mathfrak{t}_{D, P_\bullet}^{\text{Ref, HT}} / \mathfrak{t}_{D, f} \leq \sum_{\tau} \ell(\sigma_\tau),$$

Proof. Write $(\delta_1, \dots, \delta_n)$ for the parameter of P_\bullet . By Theorem 3.12 we have

$$\dim_L \mathfrak{t}_{D, P_\bullet}^{\text{Ref, HT}} / \mathfrak{t}_{D, f} \leq \sum_{1 \leq i < j \leq n} \dim_L H_{/f}^1(\delta_i \delta_j^{-1}).$$

Since the Hodge–Tate weights of D are distinct and the triangulation P_\bullet is regular generic, Lemma 3.13 above implies that if $i \neq j$, then $\dim_L H_{/f}^1(\delta_i \delta_j^{-1}) = \#\{\tau : \text{HT}_\tau(\delta_i) > \text{HT}_\tau(\delta_j)\}$. Thus Theorem 3.12 gives us

$$\begin{aligned} \dim_L \mathfrak{t}_{D, P_\bullet}^{\text{Ref, HT}} / \mathfrak{t}_{D, f} &\leq \sum_{1 \leq i < j \leq n} \dim_L H_{/f}^1(\delta_i \delta_j^{-1}) = \sum_{\tau} \#\{(i, j) : i < j \text{ and } \text{HT}_\tau(\delta_i) > \text{HT}_\tau(\delta_j)\} \\ &= \sum_{\tau} \underbrace{\#\{(i, j) : i < j \text{ and } \sigma_\tau(i) > \sigma_\tau(j)\}}_{\ell(\sigma_\tau)}. \end{aligned}$$

The final equality was the definition of the critical type (see Definition 2.2). \square

3.4. The relative tangent space for weakly-refined deformations. Suppose that D is a crystalline (φ, Γ_K) -module over $\mathcal{R}_{K,L}$ with distinct crystalline eigenvalues and distinct Hodge–Tate weights. We assume throughout this section that D is also equipped with a regular generic triangulation P_\bullet whose parameter we denote by $(\delta_1, \dots, \delta_n)$. We write $\mathfrak{t}_{D, P_\bullet}^{\text{Ref}}$ for the tangent space to the weakly-refined deformations with respect to P_\bullet .

If $\tilde{D} \in \mathfrak{t}_D$ then the uniqueness of the Hodge–Tate weights means \tilde{D} has Hodge–Sen–Tate weights $\{\eta_{i,\tau}\}_{i,\tau}$ in $L[\varepsilon]$ such that $\eta_{i,\tau} = h_{i,\tau} + \varepsilon d\eta_{i,\tau}$, where $d\eta_{i,\tau} \in L$. The map $\tilde{D} \mapsto (d\eta_{i,\tau})_{i,\tau}$ defines an L -linear map

$$\mathfrak{t}_D \xrightarrow{d\eta} \bigoplus_{\tau} L^{\oplus n}$$

and by definition of Hodge–Tate weights we have an exact sequence

$$(12) \quad 0 \rightarrow \mathfrak{t}_{D,P_{\bullet}}^{\text{Ref,HT}} / \mathfrak{t}_{D,f} \rightarrow \mathfrak{t}_{D,P_{\bullet}}^{\text{Ref}} / \mathfrak{t}_{D,f} \rightarrow \bigoplus_{\tau} L^{\oplus n}$$

of L -vector spaces.

Lemma 3.15. *Let $(\sigma_{\tau})_{\tau}$ be the critical type of P_{\bullet} . The image of $d\eta$ is contained in the subspace $\bigoplus_{\tau} V_{\sigma_{\tau}} \subset \bigoplus_{\tau} L^{\oplus n}$ where*

$$V_{\sigma_{\tau}} := \{(v_i) \in L^{\oplus n} : v_{\sigma_{\tau}(i)} = v_i \text{ for } i = 1, \dots, n\}.$$

Proof. The lemma states $\eta_{\sigma_{\tau}(i),\tau} - \eta_{i,\tau}$ is constant for each i, τ . This follows from [9, Lemma 7.2] (compare with the proofs of Lemma 3.9 and [9, Theorem 7.1]). \square

Recall that if $\sigma \in S_n$ then we write $c(\sigma)$ for the number of orbits in $\{1, \dots, n\}$ under the action of the cyclic group generated by σ .

Theorem 3.16. *Suppose that D is a crystalline (φ, Γ_K) -module over $\mathcal{R}_{K,L}$ equipped with a regular generic triangulation P_{\bullet} with critical type $(\sigma_{\tau})_{\tau}$. Then*

$$\dim \mathfrak{t}_{D,P_{\bullet}}^{\text{Ref}} / \mathfrak{t}_{D,f} \leq \sum_{\tau} \ell(\sigma_{\tau}) + c(\sigma_{\tau}).$$

Proof. It is easy to see that $\dim_L V_{\sigma_{\tau}} = c(\sigma_{\tau})$ for each embedding τ . By Lemma 3.15 and Corollary 3.14 we deduce that

$$\dim_L \mathfrak{t}_{D,P_{\bullet}}^{\text{Ref}} / \mathfrak{t}_{D,f} \leq \dim_L \mathfrak{t}_{D,P_{\bullet}}^{\text{Ref,HT}} / \mathfrak{t}_{D,f} + \sum_{\tau} c(\sigma_{\tau}) \leq \sum_{\tau} \ell(\sigma_{\tau}) + c(\sigma_{\tau}),$$

as we wanted. \square

To end this section we briefly explain the upper bound we have produced. If $\sigma \in S_n$ then let $\text{ord}(\sigma)$ be its order as an element of S_n . We leave the following lemma for the reader.

Lemma 3.17. *If $\sigma \in S_n$ is a cycle then $\ell(\sigma) + 1 \geq \text{ord}(\sigma)$ with equality if and only if σ is a product of distinct simple transpositions.*

Proposition 3.18. *Let $\sigma \in S_n$. Then $\ell(\sigma) + c(\sigma) \geq n$ with equality if and only if σ is a product of distinct simple transpositions.*

Proof. Write $\sigma = \sigma_1 \cdots \sigma_r$ where the σ_i are disjoint cycles. In particular, σ is a product of distinct simple transpositions if and only if each σ_i is. Next, we note that

- $\ell(\sigma) = \sum \ell(\sigma_i)$ and
- $c(\sigma) = n + r - \sum \text{ord}(\sigma_i)$.

Thus we have

$$\ell(\sigma) + c(\sigma) = n + \sum_{i=1}^r (\ell(\sigma_i) + 1 - \text{ord}(\sigma_i)).$$

Lemma 3.17 implies that the terms in the sum are all non-negative. This shows $\ell(\sigma) + c(\sigma) \geq n$ always. Moreover, we have equality if and only if $\ell(\sigma_i) + 1 - \text{ord}(\sigma_i) = 0$ for all $i = 1, \dots, r$. But Lemma 3.17 also implies that this is equivalent to σ_i being a product of distinct simple transpositions for each $i = 1, \dots, r$, so we are done. \square

Corollary 3.19. *Suppose that D is a crystalline (φ, Γ_K) -module over $\mathcal{R}_{K,L}$ equipped with a regular generic triangulation P_\bullet with critical type $(\sigma_\tau)_\tau$. If each σ_τ is a product of distinct simple transpositions then $\dim_L \mathfrak{t}_{D,P_\bullet}^{\text{Ref}} / \mathfrak{t}_{D,f} \leq (K : \mathbf{Q}_p) \cdot n$.*

Proof. Combine Proposition 3.18 and Theorem 3.16. \square

Remark 3.20. Ostroff showed the author an easy argument that the number of $\sigma \in S_n$ which are products of distinct simple transpositions is given by F_{2n} where F_m is the Fibonacci sequence starting with $F_1 = 0$ and $F_2 = 1$. In particular, the proportion $F_{2n}/n!$ of permutations which are products of distinct simple transpositions tends to zero as $n \rightarrow +\infty$.

4. APPLICATION TO EIGENVARIETIES

4.1. Unitary groups and Galois representations. Our goal in this subsection is to specify notations and conventions for unitary groups, automorphic representations, and Galois representations. We do not strive for the greatest generality; our goal is to illustrate how the local deformation calculation in Section 3 can be used to bound dimensions of tangent spaces on eigenvarieties. Our hypotheses may be weakened in various directions, especially as progress is made in constructing Galois representations and Langlands functoriality.

Let F/F^+ be a CM extension of number fields with F^+ totally real and F a totally imaginary quadratic extension of F^+ . We assume F/F^+ is unramified everywhere and each p -adic place of F^+ splits in F .

Fix an integer $n \geq 1$ which is either odd or if n is even then assume that $n(F^+ : \mathbf{Q}) \equiv 0 \pmod{4}$. With this, we let \mathbf{G} denote a unitary group in n variables over F^+ such that

- \mathbf{G} is split over F . We fix an isomorphism $\mathbf{G} \times_{F^+} F \simeq \text{GL}_{n/F}$.
- \mathbf{G} is quasi-split at each finite place of F^+ .
- $\mathbf{G}(F^+ \otimes_{\mathbf{Q}} \mathbf{R})$ is a finite product of copies of the compact real unitary group $\text{U}(n)$.

We refer to \mathbf{G} as a definite unitary group associated to F/F^+ .

If w is a place of F^+ then let F_w^+ denote the corresponding local field (and similarly for places of F). If w splits in F then the choice of $\tilde{w} \mid w$ determines an isomorphism $\mathbf{G}(F_w^+) \simeq \text{GL}_n(F_{\tilde{w}}) = \text{GL}_n(F_w^+)$ which implicitly depends on \tilde{w} .

We fix a compact open subgroup $U^p = \prod_{w \nmid p} U_w \subset \mathbf{G}(\mathbf{A}_{F^+}^{p\infty})$. Here, $\mathbf{A}_{F^+}^{p\infty}$ is the finite adeles of F^+ away from p . We assume throughout that U_w is maximal hyperspecial compact at every inert place of F^+ . Finally, we also write S for a finite set of finite places of F^+ such that S contains all the p -adic places and all the places w such that U_w is not maximal hyperspecial compact. In particular, each place in S is split in F . We will also use S for the set of places \tilde{w} of F such that $\tilde{w} \mid w$ with $w \in S$.

Choose an isomorphism $\mathbf{C} \simeq \overline{\mathbf{Q}}_p$. Then, each complex embedding $v_\infty : F^+ \hookrightarrow \mathbf{C}$ corresponds uniquely to a pair (v, τ) where v is a p -adic place of F^+ and $\tau : F_v^+ \hookrightarrow \overline{\mathbf{Q}}_p$ is an embedding. Given the pair (v, τ) we write $(v, \tau)_\infty$ for the corresponding infinite place of F^+ .

Automorphic representations are irreducible direct summands in the space of complex-valued functions on $\mathbf{G}(F^+) \backslash \mathbf{G}(\mathbf{A}_{F^+})$ which are smooth and $G(F^+ \otimes_{\mathbf{Q}} \mathbf{R})$ -finite (see [5, Section

6.2.3] for example). Every automorphic representation π may be factored as $\pi = \pi_\infty \otimes \pi_f$, where $\pi_f = \bigotimes' \pi_w$ is a representation of $\mathbf{G}(\mathbf{A}_{F^+}^\infty)$ and π_∞ is the weight of π , which is an algebraic representation of $\mathbf{G}(F^+ \otimes_{\mathbf{Q}} \mathbf{R})$. The weight π_∞ factors as a tensor product $\bigotimes_{v_\infty} \pi_{v_\infty}$ of irreducible algebraic representations of $\mathrm{U}(n)$ indexed by infinite places v_∞ of F^+ . Let $k_{v_\infty} = (k_{1,v_\infty} \geq k_{2,v_\infty} \geq \cdots \geq k_{n,v_\infty})$ be the dominant weight associated to π_{v_∞} . If (v, τ) is a pair as in the previous paragraph we write $k_{i,v,\tau} := k_{i,(v,\tau)_\infty}$ and for fixed v, τ we stress that $k_{1,v,\tau} \geq k_{2,v,\tau} \geq \cdots \geq k_{n,v,\tau}$. We say that π has tame level U^p if $\pi_f^{U^p} \neq 0$.

By the work of many authors, for each automorphic representation π of tame level U^p there is a unique n -dimensional continuous semi-simple representation $\rho_\pi : G_{F,S} \rightarrow \mathrm{GL}_n(\overline{\mathbf{Q}}_p)$ such that:⁸

- (LCG-I) ρ_π conjugate self-dual up to a twist, i.e. $\rho_\pi^\perp \simeq \rho_\pi(n-1)$ where ρ^\perp is the conjugate dual representation $g \mapsto {}^t \rho(\tilde{c}g\tilde{c})^{-1}$ ($\tilde{c} \in G_{F^+}$ is any order two lift of the non-trivial element in $\mathrm{Gal}(F/F^+)$; see [6]).
- (LCG-II) If $w \nmid p$ is a place of F^+ and $\tilde{w} \mid w$ is a place of F then $\mathrm{WD}(\rho_{\pi,\tilde{w}}) = \mathrm{rec}(\pi_{\tilde{w}}^{\tilde{w}} |\det|^{\frac{1-n}{2}})$, where $\pi_{\tilde{w}}^{\tilde{w}}$ is either the irreducible smooth representation of $\mathrm{GL}_n(F_{\tilde{w}})$ associated to π_w via the isomorphism $\mathbf{G}(F_w^+) \simeq \mathrm{GL}_n(F_{\tilde{w}})$ if w is split in F , or $\pi_{\tilde{w}}^{\tilde{w}}$ is the base change (see [37], for example) from $G(F_w^+)$ to $\mathrm{GL}_n(F_{\tilde{w}})$ of the necessarily unramified representation π_w of $G(F_w^+)$ if w is inert (and thus $w \notin S$).
- (LCG-III) If v is a p -adic place of F^+ and $\tilde{v} \mid v$ is a place of F then the representation $\rho_{\pi,\tilde{v}}$ is potentially semi-stable and $\mathrm{WD}(\rho_{\pi,\tilde{v}}) = \mathrm{rec}(\pi_{\tilde{v}}^{\tilde{v}} |\det|^{\frac{1-n}{2}})$ (with the notation as above). If π_v is unramified then $\rho_{\pi,\tilde{v}}$ is crystalline.

The Hodge–Tate weights are as follows. If $\tau : F_{\tilde{v}} \hookrightarrow \overline{\mathbf{Q}}_p$ is an embedding then the natural equality $F_v^+ = F_{\tilde{v}}$ defines a pair (v, τ) of a p -adic place v of F^+ together with an embedding $\tau : F_v^+ \hookrightarrow \overline{\mathbf{Q}}_p$. Then, the τ -Hodge–Tate weights of $\rho_{\pi,\tilde{v}}$ are given by $h_{i,\tilde{v},\tau} = -k_{i,v,\tau} + i - 1$.

Generally, the representations ρ_π are constructed in two steps. The first, requiring that \mathbf{G} is quasi-split at each finite place, is to apply the base change theorems of Labesse to [32] to transfer to automorphic representations for $\mathrm{GL}_{n/F}$. (We used that F/F^+ is everywhere unramified to not have to address ramified primes in (LCG-II).)

The second step is the vast collection of works on constructing Galois representations for regular algebraic essentially conjugate self-dual representations of $\mathrm{GL}_{n/F}$ along with their local properties. See [40, 23] for further references (along with [2, 18, 19] for the various compatibilities, especially the compatibility at $w \in S$ with $w \nmid p$ given by [18]).

4.2. Refinements and eigenvarieties. An eigenvariety p -adically interpolates automorphic representations π for \mathbf{G} , together with triangulations of the corresponding crystalline Galois representation (or orderings of crystalline eigenvalues).

Let us be more precise. Write $\mathcal{H}(U^p)^{\mathrm{sph}}$ for the spherical Hecke algebra of tame level U^p . For the next two paragraphs, fix an automorphic representation π for \mathbf{G} of tame level U^p . Then, π naturally gives rise to a ring homomorphism $\lambda_\pi : \mathcal{H}(U^p)^{\mathrm{sph}} \rightarrow \overline{\mathbf{Q}}_p$. By (LCG-II) and the Chebotarev density theorem, ρ_π is determined by λ_π .

⁸The notations and conventions may be found in Section 1.5. Below, if \tilde{w} is a place of F , we write $\rho_{\pi,\tilde{w}}$ for the restriction of ρ_π to a decomposition group of \tilde{w} . Everything depends the isomorphism $\mathbf{C} \simeq \mathbf{Q}_p$.

For each p -adic place v of F^+ we now *choose* a distinguished place $\tilde{v} \mid v$ in F .⁹ Thus we have an identification $\mathbf{G}(F_v^+) = \mathrm{GL}_n(F_v^+)$ that remains set throughout the rest of this section. We let $T(F_v^+)$, respectively $B(F_v^+)$, denote the subgroups of $\mathbf{G}(F_v^+)$ corresponding to the diagonal matrices, respectively the Borel subgroup of upper triangular matrices, and also the actual subgroups of $\mathrm{GL}_n(F_v^+)$.

If v is a p -adic place of F^+ and π_v is unramified then we may also *choose* a smooth unramified character $\vartheta_{\tilde{v}} : T(F_v^+) \rightarrow \overline{\mathbf{Q}}_p^\times$ such that $\pi_v \hookrightarrow \mathrm{Ind}_{B(F_v^+)}^{\mathrm{GL}_n(F_v^+)}(\delta_{B(F_v^+)}^{1/2} \vartheta_{\tilde{v}})$ (recall the notations from Section 1.5). Note that $\vartheta_{\tilde{v}}$ actually depends on \tilde{v} , as the identification of $\mathrm{GL}_n(F_v^+)$ with $\mathbf{G}(F_v^+)$ does. Following [5, Chapter 6], we call $\vartheta_{\tilde{v}}$ an accessible refinement for π_v . We also denote by $\psi_{\tilde{v}} : T(F_v^+) \rightarrow \overline{\mathbf{Q}}_p^\times$ the highest weight of the unique irreducible algebraic representation $\mathrm{Res}_{F_v^+/\mathbf{Q}_p} \mathrm{GL}_n$ over $\overline{\mathbf{Q}}_p$ with weight given by the tuple $(k_{i,v,\tau})_{i,\tau}$. Explicitly, if $z \in (F_v^+)^\times$ then $\psi_{\tilde{v}}(\mathrm{diag}(1, 1, \dots, z, \dots, 1, 1)) = \prod_{\tau} \tau(z)^{k_{i,v,\tau}}$ where z appears in the i th spot on the left-hand side of that equation.

Now let $T = \prod_v T(F_v^+)$ and \widehat{T} be the rigid analytic space over \mathbf{Q}_p parameterizing continuous character of T . Then, for each automorphic representation π of tame level U^p which is unramified at the p -adic places we have a point $\chi_\pi := (\delta_{B(F_v^+)}^{-1/2} \psi_{\tilde{v}} \vartheta_{\tilde{v}})_{v|p}$ in \widehat{T} , depending on the choice of accessible refinements $\vartheta_{\tilde{v}}$. The eigenvariety X_{U^p} of tame level U^p is coarsely defined as the rigid analytic closure of the points

$$Z_{\mathrm{cl}} := \{(\lambda_\pi, \chi_\pi)\} \subset \mathrm{Hom}(\mathcal{H}(U^p)^{\mathrm{sph}}, \overline{\mathbf{Q}}_p) \times \widehat{T}$$

where the collection runs over automorphic representations π of tame level U^p , unramified at the p -adic places, together with the choice of an accessible refinement at each p -adic places. We call Z_{cl} the set of “classical points”. The rigid analytic closure does not literally make sense, but we refer to [20, 24] for details and precisions on the construction. For the remainder of this section, we summarize the properties that we will need.

The natural map $\chi : X_{U^p} \rightarrow \widehat{T}$ is written $x \mapsto \chi_x$. By way of comparison with other sources, the map χ contains the data that may usually be included in the presence of the Atkin–Lehner algebra (in the style of [20, 5] for example).

We briefly observe the role of our normalizations. If (λ_π, χ_π) is a classical point, and if $v \mid p$ is a p -adic place we write $\chi_{1,v,\pi} \otimes \dots \otimes \chi_{n,v,\pi} : T(F_v^+) \rightarrow \overline{\mathbf{Q}}_p^\times$ for the local component at v of χ_π . Then, (LCG-III) implies that the crystalline eigenvalues of $\rho_{\pi,\tilde{v}}$ are given by the list

$$(13) \quad \left(\chi_{1,v,\pi}(\varpi_v) \cdot \prod_{\tau} \tau(\varpi_v)^{h_{1,v,\tau}}, \dots, \chi_{n,v,\pi}(\varpi_v) \cdot \prod_{\tau} \tau(\varpi_v)^{h_{n,v,\tau}} \right)$$

for some/any choice of uniformizer $\varpi_v \in F_v^+$. In particular, each classical point (with distinct crystalline eigenvalues) is naturally equipped with triangulations at the p -adic places.

The classical points Z_{cl} are Zariski dense and accumulating in X_{U^p} [5, Theorem 7.3.1(v)]. The p -adic analytic variation of pseudocharacters may be used to construct a global pseudocharacter $T : G_{F,S} \rightarrow \mathrm{GL}_n(\mathcal{O}(X_{U^p}))$ which interpolates $x \mapsto \mathrm{tr}(\rho_x)$ at classical points [20, Proposition 7.1.1]. Specializing T to a point x , [43, Theorem 1(2)] also gives a continuous semi-simple representation

$$\rho_x : G_{F,S} \rightarrow \mathrm{GL}_n(\overline{\mathbf{Q}}_p)$$

⁹This choice is ultimately inconsequential. See the remark preceding Proposition 4.5.

which is conjugate self-dual and satisfies apparent compatibility over X_{U^p} at unramified places (by interpolation). If $x \in X$ we let $L(x)$ denote its residue field.

Lemma 4.1. *If $x \in X_{U^p}$ and ρ_x is absolutely irreducible and defined over $L(x)$ then there exists an affinoid neighborhood $Y = \mathrm{Sp}(B) \subset X$ and a continuous representation $\rho_Y : G_{F,S} \rightarrow \mathrm{GL}_n(B)$ such that $\rho_Y \otimes_B L(y) = \rho_y$ for all $y \in Y$.*

Proof. Let A be the rigid local ring $A = \mathcal{O}_{X,x}^{\mathrm{rig}}$. Since A is Henselian ([12, Theorem 2.1.5]) and ρ_x is absolutely irreducible, ρ_x lifts uniquely to a continuous representation $\rho_A : G_{F,S} \rightarrow \mathrm{GL}_n(A)$ [39, Corollarie 5.2]. The existence of $Y = \mathrm{Sp}(B)$ and ρ_Y such that $\mathrm{tr}(\rho_y) = \mathrm{tr}(\rho_Y \otimes_B L(y))$ for all $y \in Y$ follows from [5, Lemma 4.3.7]. But, the locus of points $y \in Y$ for which ρ_y is absolutely irreducible is a rigid open subspace ([20, Section 7.2.1]), so we can shrink Y and assume that ρ_y is absolutely irreducible for all y . In that case the equality $\mathrm{tr}(\rho_y) = \mathrm{tr}(\rho_Y \otimes_B L(y))$ implies that $\rho_y = \rho_Y \otimes_B L(y)$, finishing the proof. \square

Remark 4.2. The representations ρ_x for classical x satisfy the hypothesis of the lemma, i.e. they are defined over their residue fields.

Instead of working with the whole tame level U^p eigenvariety, we will instead consider a minimal eigenvariety. We briefly explain, but see [5, Example 7.5.1] for further information (see also [21, Section 3.6]). Fix an automorphic representation π , unramified at the p -adic places, and the choice of an accessible refinement giving rise to a point $x_\pi \in X_{U^p}$. The places $w \in S$ with $w \nmid p$ are all split. Fix a choice $\tilde{w} \mid w$ for each such w . Then the representation $\pi_{\tilde{w}}^{\tilde{w}}$ of $\mathbf{G}(F_w^+) \simeq \mathrm{GL}_n(F_w^+)$ has a K -type ([5, Section 6.5]), which gives rise to idempotents e_w (independent of \tilde{w}) commuting with $\mathcal{H}(U^p)^{\mathrm{sph}}$ inside the space of compactly supported continuous complex-valued functions $\mathcal{C}_c^0(\mathbf{G}(\mathbf{A}_{F^+}^p))$. The minimal eigenvariety X for $x = x_\pi$ is then the idempotent-type eigenvariety ([5, Section 7.3]) obtained from the idempotents $(e_w)_{w \in S, w \nmid p}$. This defines a closed rigid subvariety $X \hookrightarrow X_{U^p}$. The corresponding classical points are those $(\lambda_\pi, \chi_\pi) \in Z_{\mathrm{cl}}$ above such that $e_w(\pi) \neq 0$ for $w \in S$, $w \nmid p$.

We will need a property of X relating to Galois representations at the ramified places S . Let $w \in S$ and \tilde{w} be a place of F above w . If (r, N) and (r', N') are two Weil–Deligne representations of the Weil group $W_{F_{\tilde{w}}}$ then we will use the notation $N \prec_{\tilde{w}} N'$ for the “less monodromy” notation $N \prec_{I_{\tilde{w}}} N'$ introduced in [5, Definition 7.8.18] and $N \sim_{\tilde{w}} N'$ for the obvious equal version. We note two things:

- (i) If $N \sim_{\tilde{w}} N'$ then $r|_{I_{\tilde{w}}} \simeq r'|_{I_{\tilde{w}}}$. This is by definition of the relation \prec .
- (ii) Let $x \in X_{U^p}$ be classical and X be its minimal eigenvariety. If z is a classical point on X then $N_{z,\tilde{w}} \prec_{\tilde{w}} N_{x,\tilde{w}}$ for all $w \in S$, where $(r_{z,\tilde{w}}, N_{z,\tilde{w}}) = \mathrm{WD}(\rho_{z,\tilde{w}})$ is the Weil–Deligne representation associated to $\rho_{z,\tilde{w}}$. This follows from (LCG-II) and the definition of the idempotents e_w (compare with [5, Section 6.5]).

We now summarize the rest of the properties of the minimal eigenvariety:

Proposition 4.3. *Let $x \in X_{U^p}$ be a classical point and X its minimal eigenvariety.*

- (a) X is equidimensional of dimension $(F^+ : \mathbf{Q}) \cdot n$.
- (b) If ρ_x is absolutely irreducible then there exists a canonical lifting $\hat{\rho}_x$ to $\mathcal{O}_{X,x}^{\mathrm{rig}}$, and for each $\tilde{w} \mid w$ with $w \in S$ and $w \nmid p$, we have $\hat{\rho}_x|_{I_{\tilde{w}}} \simeq \rho_x|_{I_{\tilde{w}}} \otimes_{L(x)} \mathcal{O}_{X,x}^{\mathrm{rig}}$.

(c) If ρ_x is absolutely irreducible, $\widehat{\rho}_x$ is as in part (b), and \mathcal{H}_x denotes the image of $\mathcal{H}(U^p)^{\text{sph}}$ in the local ring $\mathcal{O}_{X,x}^{\text{rig}}$ then \mathcal{H}_x is contained in the sub-algebra of $\mathcal{O}_{X,x}^{\text{rig}}$ generated by $\text{tr } \wedge^i \widehat{\rho}_x(G_{F,S})$ for $i = 1, 2, \dots, n$.

Proof. The statement (a) is a general property of eigenvarieties of idempotent-type attached to definite unitary groups [5, Theorem 7.3.1(a)].

The lifting in (b) is deduced from Lemma 4.1. The constancy of inertia acting is deduced from [5, Corollary 7.5.10]. This is explained in the proof of [5, Proposition 7.6.10], but since it is where we use the minimal eigenvariety (specifically comment (ii) above), we include the argument for convenience. Let \mathcal{K}_x be the total ring of fractions of $\mathcal{O}_{X,x}^{\text{rig}}$ and write $\mathcal{K}_x = \prod \mathcal{K}_{s(x)}$ where each $\mathcal{K}_{s(x)}$ is a field corresponding to an irreducible component $s(x)$. Let $\rho_x^{\text{gen}} : G_{F,S} \rightarrow \text{GL}_n(\mathcal{K}_{s(x)})$ be the corresponding Galois representation and $\rho_{s(x),\widetilde{w}}^{\text{gen}}$ be its restriction to the local group $G_{F_{\widetilde{w}}}$. This representation admits a Weil–Deligne representation $(r_{s(x),\widetilde{w}}^{\text{gen}}, N_{s(x),\widetilde{w}}^{\text{gen}})$ (see [5, Section 7.8.4]). In general, $N_{z,\widetilde{w}} \prec_{\widetilde{w}} N_{s(x),\widetilde{w}}^{\text{gen}}$ for all z on $s(x)$ with $\sim_{\widetilde{w}}$ on a Zariski-dense subset. For instance, $N_{z,\widetilde{w}} \sim_{\widetilde{w}} N_{s(x),\widetilde{w}}^{\text{gen}}$ for a set of classical z on $s(x)$ accumulating at x . Thus (ii) above implies $N_{s(x),\widetilde{w}}^{\text{gen}} \prec_{\widetilde{w}} N_{x,\widetilde{w}}$. The reverse is always true, so we conclude $N_{s(x),\widetilde{w}}^{\text{gen}} \sim_{\widetilde{w}} N_{x,\widetilde{w}}$. Since $s(x)$ is arbitrary, we see that the hypothesis of [5, Corollary 7.5.10] is satisfied.¹⁰ We conclude that $\widehat{N}_{x,\widetilde{w}} \sim_{\widetilde{w}} N_{x,\widetilde{w}}$ where $(\widehat{r}_{x,\widetilde{w}}, \widehat{N}_{x,\widetilde{w}})$ is the Weil–Deligne representation associated to $\widehat{\rho}_{x,\widetilde{w}}$ (see the references just prior to [5, Corollary 7.5.10]). By definition then $\widehat{\rho}_x|_{F_{\widetilde{w}}} \simeq \rho_x|_{F_{\widetilde{w}}} \otimes_{L(x)} \mathcal{O}_{X,x}^{\text{rig}}$ (see (i) above).

Part (c) follows from applying (LCG-II) to the places $w \notin S$. Indeed, the Satake isomorphism and (LCG-II) implies that $\mathcal{H}(U^p)^{\text{sph}}$ is generated as an algebra by elements whose specialization at any classical point z are the coefficients of the characteristic polynomial of $\rho_{z,\widetilde{w}}(\text{Frob}_{\widetilde{w}})$. In particular, the image \mathcal{H}_x in $\mathcal{O}_{X,x}^{\text{rig}}$ is naturally generated by the characteristic polynomials of $\widehat{\rho}_x(\text{Frob}_w)$ with $w \notin S$. \square

4.3. Upper bounds for tangent spaces via deformation theory. We continue to use the notations and conventions of the previous two sections. We also assume:

(4.3-A) π is an automorphic representation of tame level U^p , unramified above p , and $(\vartheta_{\widetilde{v}})_{\widetilde{v}}$ is a list of accessible refinements for the representations π_v at the p -adic places v of F^+ .

(4.3-B) The global Galois representation ρ_π is absolutely irreducible.

(4.3-C) By (LCG-III) the choices in (4.3-A) define orderings of the crystalline eigenvalues for each $\rho_{\pi,\widetilde{v}}$ (see (13)). We *assume* that these eigenvalues are all distinct and denote by $P_{\widetilde{v},\bullet}$ the corresponding triangulation of the $(\varphi, \Gamma_{F_v^+})$ -module $D_{\text{rig}}^\dagger(\rho_{\pi,\widetilde{v}})$.

(4.3-D) We *further* assume that $P_{\widetilde{v},\bullet}$ is regular generic for each $v \mid p$.

We now denote by $X \subset X_{U^p}$ the minimal eigenvariety containing the point $x = x_\pi$. By assumption (4.3-B), ρ_x is absolutely irreducible and so the universal deformation functor \mathfrak{X}_{ρ_x} on $\mathfrak{A}_{L(x)}$ is (pro-)representable by a complete local noetherian $L(x)$ -algebra $R_{\rho_x}^{\text{univ}}$ with residue field $L(x)$. Denote by $\mathfrak{X}_{\rho_x}^{\text{csd}} \subset \mathfrak{X}_{\rho_x}$ the representable subfunctor parameterizing conjugate self-dual deformations ρ (i.e. $\rho^\perp \simeq \rho(n-1)$).

¹⁰The embedded reference to Proposition 7.5.8 in *loc. cit* is valid because we’ve assume that each place of S splits in F .

At places $\tilde{w} \nmid p$ of F , we may consider the local Galois representation $\rho_{x,\tilde{w}}$ and its universal deformation functor $\mathfrak{X}_{\rho_{x,\tilde{w}}}$. There is a natural subfunctor $\mathfrak{X}_{\rho_{x,\tilde{w}},f} \subset \mathfrak{X}_{\rho_{x,\tilde{w}}}$ parameterizing deformations which are minimally ramified. That is, if $A \in \mathfrak{A}\mathfrak{R}_{L(x)}$ then

$$\mathfrak{X}_{\rho_{x,\tilde{w}},f}(A) = \left\{ \rho_A \in \mathfrak{X}_{\rho_{x,\tilde{w}}}(A) : \rho_A|_{I_{\tilde{w}}} \simeq \rho_{x,\tilde{w}}|_{I_{\tilde{w}}} \otimes_{L(x)} A \right\}.$$

For example, if $\tilde{w} \notin S$ then $\rho_A \in \mathfrak{X}_{\rho_{x,\tilde{w}},f}(A)$ if and only if ρ_A is trivial on inertia at \tilde{w} . The arrow $\mathfrak{X}_{\rho_{x,\tilde{w}},f} \hookrightarrow \mathfrak{X}_{\rho_{x,\tilde{w}}}$ is relatively representable (e.g. by Schlessinger's criterion).

For $v \mid p$ in F^+ , we let $\mathfrak{X}_{\rho_{x,\tilde{v}}}^{\text{Ref}} := \mathfrak{X}_{D_{\text{rig}}^\dagger(\rho_{x,\tilde{v}}), P_{\tilde{v},\bullet}}^{\text{Ref}}$ be the weakly-refined deformations of $D_{\text{rig}}^\dagger(\rho_{x,\tilde{v}})$ with respect to the triangulation $P_{\tilde{v},\bullet}$ arising from (4.3-C). We now let $\mathfrak{X}_{\rho_x}^{\text{Ref,min}}$ denote the fibered product defined by the diagram

$$\begin{array}{ccc} \mathfrak{X}_{\rho_x}^{\text{Ref,min}} & \xrightarrow{\quad\quad\quad} & \mathfrak{X}_{\rho_x}^{\text{csd}} \\ \downarrow & & \downarrow \\ \prod_{\tilde{w} \nmid p} \mathfrak{X}_{\rho_{x,\tilde{w}},f} \times \prod_{v \mid p} \mathfrak{X}_{\rho_{x,\tilde{v}}}^{\text{Ref}} & \longrightarrow & \prod_{\tilde{w} \nmid p} \mathfrak{X}_{\rho_{x,\tilde{w}}} \times \prod_{v \mid p} \mathfrak{X}_{\rho_{x,\tilde{v}}}. \end{array}$$

Since the bottom arrow is relatively representable and $\mathfrak{X}_{\rho_x}^{\text{csd}}$ is itself (pro-)representable, we deduce that $\mathfrak{X}_{\rho_x}^{\text{Ref,min}}$ is a (pro-)representable functor on $\mathfrak{A}\mathfrak{R}_{L(x)}$. We denote the universal ring representing $\mathfrak{X}_{\rho_x}^{\text{Ref,min}}$ by $R_{\rho_x}^{\text{Ref,min}}$.

Remark 4.4. The conjugate self-dual property of the deformations in $\mathfrak{X}_{\rho_x}^{\text{Ref,min}}$ implies that it is sufficient to consider only one place of F above each place of F^+ . This is why, for example, we only specify the deformation problem at the fixed p -adic places \tilde{v} of F .

Let $\mathcal{O}_{X,x}^{\text{rig}}$ denote the rigid analytic local ring of x at X and $\widehat{\mathcal{O}}_{X,x}^{\text{rig}}$ denote its completion.

Proposition 4.5. *There is a canonical surjective ring homomorphism $f : R_{\rho_x}^{\text{Ref,min}} \twoheadrightarrow \widehat{\mathcal{O}}_{X,x}^{\text{rig}}$.*

Proof. Since ρ_x is absolutely irreducible, Proposition 4.3(b) implies that we can canonically lift ρ_x to a deformation $\widehat{\rho}_x : G_{F,S} \rightarrow \text{GL}_n(\mathcal{O}_{X,x}^{\text{rig}}) \subset \text{GL}_n(\widehat{\mathcal{O}}_{X,x}^{\text{rig}})$. Since $\widehat{\mathcal{O}}_{X,x}^{\text{rig}}$ is a complete local Noetherian $L(x)$ -algebra with residue field $L(x)$, this defines a map $f : R_{\rho_x}^{\text{univ}} \rightarrow \widehat{\mathcal{O}}_{X,x}^{\text{rig}}$. To show that f factors through $R_{\rho_x}^{\text{Ref,min}}$, we need to show that $\widehat{\rho}_x$ defines a point in the subfunctor $\mathfrak{X}_{\rho_x}^{\text{Ref,min}} \subset \mathfrak{X}_{\rho_x}$. For that, we need to check:

- (i) $\widehat{\rho}_x$ is conjugate self-dual.
- (ii) $\widehat{\rho}_x$ is minimally ramified away from p .
- (iii) If $v \mid p$ then the representation $\widehat{\rho}_{x,\tilde{v}}$ is a weakly-refined deformation of $\rho_{x,\tilde{v}}$.

The first point is clear because the Galois representation ρ_y is conjugate self-dual for all $y \in X$. The condition (ii) follows from Proposition 4.3(b), which is valid by assumption (4.3-B) and the minimality of X . The point (iii) is the crucial p -adic interpolation of crystalline eigenvalues over eigenvarieties as we now explain (see [31, 5, 34, 30]).

Let Y be an affinoid open of X , containing x , as in Lemma 4.1. Fix a place $v \mid p$ in F^+ and form the family $D_{\text{rig}}^\dagger(\rho_{Y,\tilde{v}})$ of (φ, Γ_K) -modules over Y (see [29]). Consider the natural map $\chi : Y \rightarrow \widehat{T}$ as the universal character $\chi : T \rightarrow \mathcal{O}(Y)^\times$.¹¹ Write $\chi_v : T(F_v^+) \rightarrow \mathcal{O}(Y)^\times$

¹¹Here and below we are using $\mathcal{O}(-)$ to denote the ring of rigid analytic functions on a rigid space.

for the local component at v , and we further write $\chi_v = \chi_{1,v} \otimes \cdots \otimes \chi_{n,v}$, with each $\chi_{j,v}$ a character of $(F_v^+)^{\times}$. Set $\Delta_{j,v} := \chi_{1,v} \cdots \chi_{j,v} : (F_v^+)^{\times} \rightarrow \mathcal{O}(Y)^{\times}$.

Choose a uniformizer $\varpi_{F_v^+} \in (F_v^+)^{\times}$ and use $\varpi_{F_v^+}$ to write $(F_v^+)^{\times} = \varpi_{F_v^+}^{\mathbb{Z}} \times \mathcal{O}_{F_v^+}^{\times}$. Let $\eta_{j,v} := \Delta_{j,v}|_{\mathcal{O}_{F_v^+}^{\times}}$, but then use the same notation to denote what we called $(\eta_{j,v})_{\varpi_K}$ in Section 2.3, i.e. set $\eta_{j,v}(\varpi_K) = 1$. With these notations, [34] implies that for $j = 1, \dots, n$ the space $D_{\text{crys}}^+(\wedge^j D_{\text{rig}}^{\dagger}(\rho_{Y,\tilde{v}})(\eta_{j,v}^{-1}))^{\varphi^{fK}=\Delta_{j,v}(\varpi_{F_v^+})}$ is a coherent sheaf of generic rank one on Y (see the comments preceding [34, Theorem 0.3.4] and note the consistency with (13)).

We can say more. By assumption (4.3-D), the triangulations $P_{v,\bullet}$ at the point x are all regular generic and thus [34, Proposition 4.3.5] implies that after shrinking Y we may assume that each $D_{\text{crys}}^+(\wedge^j D_{\text{rig}}^{\dagger}(\rho_{Y,\tilde{v}})(\eta_{j,v}^{-1}))^{\varphi^{fK}=\Delta_{j,v}(\varpi_{F_v^+})}$ is free of rank one, and satisfies base change, on Y . In particular, this shows that $\widehat{\rho}_{x,\tilde{v}}$ is a weakly-refined deformation of $\rho_{x,\tilde{v}}$.

We now give the standard argument that the map f is surjective. By compactness of the ring $R_{\rho_x}^{\text{Ref},\text{min}}$, it is enough to see that dense subring $\mathcal{O}_{X,x}^{\text{rig}} \subset \widehat{\mathcal{O}}_{X,x}^{\text{rig}}$ is contained in the image of f . By construction of X , $\mathcal{O}_{X,x}^{\text{rig}}$ is topologically generated by $\mathcal{H}(U^p)^{\text{sph}}$ over $\mathcal{O}(\widehat{T})$ inside $\mathcal{O}_{X,x}^{\text{rig}}$ (compare with the proof of [5, Proposition 7.6.10]). The image of the algebra $\mathcal{H}(U^p)^{\text{sph}}$ in $\mathcal{O}_{X,x}^{\text{rig}}$ is in the image of f by Proposition 4.3(c).

The image of $\mathcal{O}_{\widehat{T},x}^{\text{rig}}$ in $\mathcal{O}_{X,x}^{\text{rig}}$ is generated by $\{\Delta_{j,v}, \eta_{j,v}\}$ for $v \mid p$ and $j = 1, \dots, n$ as above (meaning the values of these characters as function on Y). Let ρ_x^{univ} be the universal deformation to $R_{\rho_x}^{\text{Ref},\text{min}}$. If $I \subset R_{\rho_x}^{\text{Ref},\text{min}}$ is a co-finite length ideal, we apply the definition of being weakly-refined with the choice of uniformizer $\varpi_{F_v^+}$ as above, and deduce that $D_{\text{crys}}^+(\wedge^j D_{\text{rig}}^{\dagger}(\rho_{x,\tilde{v}}^{\text{univ}}/I)(\eta_{\text{univ},j,v}^{-1}))^{\varphi^{fK}=\Delta_{\text{univ},j,v}(\varpi_{F_v^+})}$ is free of rank one over $R_{\rho_x}^{\text{Ref},\text{min}}/I \otimes_{L(x)} (F_v^+)_0$, where $\Delta_{\text{univ},j,v}$ and $\eta_{\text{univ},j,v}$ are the universal characters of $(F_v^+)^{\times} \rightarrow (R_{\rho_x}^{\text{Ref},\text{min}})^{\times}$ associated to ρ_x^{univ} . Since ρ_x^{univ} induces $\widehat{\rho}_x$ under the map f , we see that the corresponding compositions $(F_v^+)^{\times} \rightarrow (R_{\rho_x}^{\text{Ref},\text{min}}/I)^{\times} \rightarrow (\mathcal{O}_{X,x}^{\text{rig}}/I\mathcal{O}_{X,x}^{\text{rig}})^{\times}$ are equal to $\Delta_{j,v} \bmod I$ and $\eta_{j,v} \bmod I$, respectively (we are carefully using that $\eta_{\text{univ},j,v}$ and $\eta_{j,v}$ are both defined to be trivial on $\varpi_{F_v^+}$). Now we take the limit over I being powers of the maximal ideal in $\mathcal{O}_{X,x}^{\text{rig}}$ and conclude $\{\Delta_{j,v}, \eta_{j,v}\}$ are in the image of f by Krull's intersection theorem. \square

Now denote by $T_{X,x}$ the tangent space to X at the point x . Let $\mathfrak{t}_{\rho_x}^{\text{Ref},\text{min}}$ denote the tangent space for the ring $R_{\rho_x}^{\text{Ref},\text{min}}$. If $\dim_{L(x)} \mathfrak{t}_{\rho_x}^{\text{Ref},\text{min}} = g$ then $R_{\rho_x}^{\text{Ref},\text{min}}$ is a quotient of a power series ring $L(x)[[u_1, \dots, u_g]]$ in g variables (this goes back to Mazur).

Since $\rho_x^{\perp} \simeq \rho_x(n-1)$, there is a matrix $A_x \in \text{GL}_n(L(x))$ such that $\rho_x^{\perp}(g) = A_x \rho_x(g) A_x^{-1} \chi_{\text{cycl}}^{n-1}(g)$ for all $g \in G_F$. The main theorem of [6] implies that A_x is symmetric and unique up to scalar because ρ_x is absolutely irreducible. We recall that this allows us to extend the action of G_F on $\text{ad } \rho_x$ to an action of G_{F^+} (see the introduction of [1], for example). Specifically, we write $\text{ad } \rho_x = \text{GL}(M_n(L(x)))$, and if $c \in G_{F^+}$ denotes the choice of a complex conjugation for F/F^+ then we let c act by $B \mapsto -A_x^{-1} \cdot {}^t B \cdot A_x$ for all $B \in M_n(L)$. This is well-defined and completely canonical. Let $H_f^1(\text{ad } \rho_x)^+ = H_f^1(G_{F^+}, \text{ad } \rho_x)$ denote the global adjoint Bloch–Kato Selmer group [13].

Lemma 4.6. *The restriction map $\rho_x \mapsto (\rho_{x,\tilde{v}})$ induces a natural map $\mathfrak{t}_{\rho_x}^{\text{Ref},\min} \rightarrow \bigoplus_{v|p} \mathfrak{t}_{\rho_{x,\tilde{v}}}^{\text{Ref}}$ and this induces an exact sequence*

$$0 \rightarrow H_f^1(\text{ad } \rho_x)^+ \rightarrow \mathfrak{t}_{\rho_x}^{\text{Ref},\min} \rightarrow \bigoplus_{v|p} \mathfrak{t}_{\rho_{x,\tilde{v}}}^{\text{Ref}} / \mathfrak{t}_{\rho_{x,\tilde{v}},f}$$

of vector spaces over $L(x)$.

Proof. The fact that the natural map exists is clear. Next, $\text{Gal}(F/F^+) = \{1, c\}$ acts on $H^1(G_F, \text{ad } \rho_x)$ by the paragraph preceding this lemma, and the inflation-restriction sequence implies that $H^1(G_{F^+}, \text{ad } \rho_x) = H^1(G_F, \text{ad } \rho_x)^{c=1}$. It is an elementary calculation that a deformation $\tilde{\rho} \in H^1(G_F, \text{ad } \rho_x)$ of ρ_x to $L(x)[\varepsilon]$ is conjugate self-dual if and only if $\tilde{\rho} \in H^1(G_F, \text{ad } \rho_x)^{c=1}$. Thus $H_f^1(\text{ad } \rho_x)^+$ is contained in the kernel of the restriction maps. The reverse inclusion follows from the minimal condition since every deformation $\tilde{\rho} \in \mathfrak{t}_{\rho_x}^{\text{Ref},\min}$ has the property that the restriction $\tilde{\rho}_{\tilde{w}}$ to a place $\tilde{w} \nmid p$ lies in $H_f^1(\text{ad } \rho_{x,\tilde{w}}) = \mathfrak{t}_{\rho_{x,\tilde{w}},f}$ already. \square

We now summarize the situation. We have fixed a classical point x on an eigenvariety corresponding to the choices (4.3-A)-(4.3-D), and we restricted to the minimal eigenvariety X containing x . By (4.3-C) the point x comes naturally equipped with triangulations $P_{\tilde{v},\bullet}$ of the local Galois representation $D_{\text{rig}}^\dagger(\rho_{x,\tilde{v}})$ at a set of distinguished p -adic places \tilde{v} of F .

Theorem 4.7. *With the above notation and assumptions, for each p -adic place v of F^+ let $(\sigma_{v,\tau})_\tau$ be critical type of the triangulation $P_{\tilde{v},\bullet}$ at the point x . Then*

$$\dim_{L(x)} T_{X,x} \leq \dim_{L(x)} \mathfrak{t}_{\rho_x}^{\text{Ref},\min} \leq \dim_{L(x)} H_f^1(\text{ad } \rho_x)^+ + \sum_{v|p} \sum_{\tau: F_v^+ \rightarrow L(x)} \ell(\sigma_{v,\tau}) + c(\sigma_{v,\tau}).$$

Proof. The surjection $f: R_{\rho_x}^{\text{Ref},\min} \rightarrow \widehat{\mathcal{O}}_{X,x}^{\text{rig}}$ in Proposition 4.5 gives rise to a canonical injection $T_{X,x} \hookrightarrow \mathfrak{t}_{\rho_x}^{\text{Ref},\min}$. This proves the first inequality. For the second inequality, Lemma 4.6 implies that

$$(14) \quad \dim_{L(x)} \mathfrak{t}_{\rho_x}^{\text{Ref},\min} \leq \dim_{L(x)} H_f^1(\text{ad } \rho_x)^+ + \sum_{v|p} \dim_{L(x)} \mathfrak{t}_{\rho_{x,\tilde{v}}}^{\text{Ref}} / \mathfrak{t}_{\rho_{x,\tilde{v}},f}.$$

If $v \mid p$ then $P_{\tilde{v},\bullet}$ is regular generic by assumption (4.3-D) and so Theorem 3.16 implies that

$$(15) \quad \dim_{L(x)} \mathfrak{t}_{\rho_{x,\tilde{v}}}^{\text{Ref}} / \mathfrak{t}_{\rho_{x,\tilde{v}},f} \leq \sum_{\tau: F_v^+ \rightarrow L(x)} \ell(\sigma_{v,\tau}) + c(\sigma_{v,\tau}).$$

The second inequality in the statement of the theorem now follows from summing (15) over $v \mid p$ and inserting it into (14). \square

Corollary 4.8. *In the situation of Theorem 4.7, suppose that*

- (a) $\sigma_{v,\tau}$ is a product of distinct simple transpositions for all v and all τ , and
- (b) $H_f^1(\text{ad } \rho_x)^+ = (0)$.

Then X is smooth at x and the map $f: R_{\rho_x}^{\text{Ref},\min} \rightarrow \widehat{\mathcal{O}}_{X,x}^{\text{rig}}$ is an isomorphism.

Proof. By Proposition 3.18, the assumption (a) implies that for each v ,

$$\sum_{\tau: F_v^+ \rightarrow L(x)} \ell(\sigma_{v,\tau}) + c(\sigma_{v,\tau}) = \sum_{\tau: F_v^+ \rightarrow L(x)} n = (F_v^+ : \mathbf{Q}_p) \cdot n.$$

Thus Theorem 4.7 implies that

$$\dim_{L(x)} T_{X,x} \leq \dim_{L(x)} \mathfrak{t}_{\rho_x}^{\text{Ref},\min} \leq \sum_{v|p} (F_v^+ : \mathbf{Q}_p) \cdot n = (F^+ : \mathbf{Q}) \cdot n.$$

Let $g = (F^+ : \mathbf{Q}) \cdot n$. Then $R_{\rho_x}^{\text{Ref},\min}$ is a quotient of a power series ring over $L(x)$ in g variables. On the other hand, $\widehat{\mathcal{O}}_{X,x}^{\text{rig}}$ is equidimensional of dimension g by Proposition 4.3(a). Krull's Hauptidealsatz then implies that the surjection $f : R_{\rho_x}^{\text{Ref},\min} \twoheadrightarrow \widehat{\mathcal{O}}_{X,x}^{\text{rig}}$ is an isomorphism and that $R_{\rho_x}^{\text{Ref},\min}$ and $\widehat{\mathcal{O}}_{X,x}^{\text{rig}}$ are both power series rings in g variables. \square

We finish with a number of remarks.

Remark 4.9. The technical hypothesis on the vanishing of the Selmer group in Corollary 4.8 is expected to always hold. The current status is discussed following Corollary 1.3.

Remark 4.10. The irreducibility of the global Galois representation ρ_x plays a role, even if under-emphasized, in the proof of Theorem 4.7. Indeed, if we only take as input the inequality $\dim_{L(x)} T_{X,x} \leq \dim_{L(x)} H_f^1(\text{ad } \rho_x)^+ + \sum_{v|p} \cdots$ in Theorem 4.7, and the assumptions (a) and (b) in Corollary 4.8, then we would still be able to prove that $\dim_{L(x)} T_{X,x} = (F^+ : \mathbf{Q}) \cdot n$, meaning X would be smooth at x . But, in [3], Bellaïche has given an example of a classical point on an eigenvariety whose critical type is a 3-cycle (hence a product of distinct simple transpositions) and which is a singular point on every irreducible component it lies on. But, ρ_x is a direct sum of two representations in this case.

Remark 4.11. In [15], Breuil, Hellmann and Schraen have shown that Corollary 4.8 is optimal in the sense that X is *singular* at classical points whose critical types are not all products of distinct simple transpositions. (They had previously shown this in [16, Section 5] assuming certain modularity conjectures.)

Remark 4.12. The astute reader may have noticed that the classicality hypothesis in Theorem 4.7 may be relaxed. Surely we used classicality to know that the local representations at the p -adic places were crystalline. But, we also used it in writing the sum in Theorem 4.7 over the p -adic places in terms of the *critical type*. Even the basic idea of the critical type requires an *a priori* reasonable ordering of Hodge–Tate weights. This ordering arises in the classical setting as the corresponding dominant weight.

But, Theorem 4.7 holds also for (sufficiently generic) crystalline points (with globally irreducible Galois representations) on the eigenvarieties, provided we replace the sum $\sum_{v|p} \sum_{\tau} \cdots$ by a sum related to the Bloch–Kato Selmer dimensions as in Theorem 3.12, i.e. a dimension depending on the parameter of the corresponding point on the eigenvariety. Doing this, it seems likely that the corresponding bounds will *not* always be tight.

The easiest examples we have in mind are *companion points* on the Coleman–Mazur eigencurve. For concreteness, we may consider an overconvergent p -adic cuspform g of negative weight $2 - k$, such that $\theta^{k-1}(g)$ is the critical p -stabilization of a p -ordinary CM form of weight k . The analog of Theorem 4.7 only produces an upper bound of two for the size of the tangent space to the one-dimensional eigencurve at the point corresponding to g .

However, we learned from Bellaïche (in a preprint which is no longer publicly available) that if one could prove that g lies on a union of CM components then in fact g lies on a unique component and is smooth. We can sketch our own proof as well. If g lies on a union of CM components then the infinitesimal deformations of the Galois representation on the

eigencurve would all be locally split at p . The locally split condition is certainly not implied by the weakly-refined condition and thus imposing it would bring the bound of two which Theorem 4.7 gives down to a bound of one, which proves g is a smooth point. It bears mentioning that we do not know whether or not such forms g must *a priori* lie on a union of CM components, but it would be remarkable if that condition was *never* satisfied.

REFERENCES

- [1] P. B. Allen. Deformations of polarized automorphic Galois representations and adjoint Selmer groups. *Duke Math. J.*, 165(13):2407–2460, 2016.
- [2] T. Barnet-Lamb, T. Gee, D. Geraghty, and R. Taylor. Local-global compatibility for $l = p$, II. *Ann. Sci. École Norm. Sup. (4)*, 2012. to appear.
- [3] J. Bellaïche. Nonsmooth classical points on eigenvarieties. *Duke Math. J.*, 145(1):71–90, 2008.
- [4] J. Bellaïche. Critical p -adic L -functions. *Invent. Math.*, 189(1):1–60, 2012.
- [5] J. Bellaïche and G. Chenevier. Families of Galois representations and Selmer groups. *Astérisque*, 324:xii+314, 2009.
- [6] J. Bellaïche and G. Chenevier. The sign of Galois representations attached to automorphic forms for unitary groups. *Compos. Math.*, 147(5):1337–1352, 2011.
- [7] D. Benois. A generalization of Greenberg’s \mathcal{L} -invariant. *Amer. J. Math.*, 133(6):1573–1632, 2011.
- [8] J. Bergdall. *On the variation of (φ, Γ) -modules over p -adic families of automorphic forms*. PhD thesis, Brandeis University, 2013.
- [9] J. Bergdall. Paraboline variation over p -adic families of (φ, Γ) -modules. *Compositio Math.*, 153(1):132–174, 2017.
- [10] L. Berger. Représentations p -adiques et équations différentielles. *Invent. Math.*, 148(2):219–284, 2002.
- [11] L. Berger. Équations différentielles p -adiques et (ϕ, N) -modules filtrés. *Astérisque*, (319):13–38, 2008. Représentations p -adiques de groupes p -adiques. I. Représentations galoisiennes et (ϕ, Γ) -modules.
- [12] V. G. Berkovich. Étale cohomology for non-Archimedean analytic spaces. *Inst. Hautes Études Sci. Publ. Math.*, (78):5–161 (1994), 1993.
- [13] S. Bloch and K. Kato. L -functions and Tamagawa numbers of motives. In *The Grothendieck Festschrift, Vol. I*, volume 86 of *Progr. Math.*, pages 333–400. Birkhäuser Boston, Boston, MA, 1990.
- [14] C. Breuil. Vers le socle localement analytique pour GL_n II. *Math. Ann.*, 361(3-4):741–785, 2015.
- [15] C. Breuil, E. Hellmann, and B. Schraen. A local model for the trianguline variety and applications. *Preprint*, 2017. Available at [arXiv:1702.02192](https://arxiv.org/abs/1702.02192).
- [16] C. Breuil, E. Hellmann, and B. Schraen. Smoothness and classicality on eigenvarieties. *Invent. Math.*, 209(1):197–274, 2017.
- [17] C. Breuil, E. Hellmann, and B. Schraen. Une interprétation modulaire de la variété trianguline. *Math. Ann.*, 367(3-4):1587–1645, 2017.
- [18] A. Caraiani. Local-global compatibility and the action of monodromy on nearby cycles. *Duke Math. J.*, 161(12):2311–2413, 2012.
- [19] A. Caraiani. Monodromy and local-global compatibility for $l = p$. *Algebra Number Theory*, 8(7):1597–1646, 2014.
- [20] G. Chenevier. Familles p -adiques de formes automorphes pour GL_n . *J. Reine Angew. Math.*, 570:143–217, 2004.
- [21] G. Chenevier. Une application des variétés de Hecke des groupes unitaires. *Preprint*, 2009.
- [22] G. Chenevier. On the infinite fern of Galois representations of unitary type. *Ann. Sci. Éc. Norm. Supér. (4)*, 44(6):963–1019, 2011.
- [23] G. Chenevier and M. Harris. Construction of automorphic Galois representations, II. *Cambridge Journal of Mathematics*, 1(1), 2013.
- [24] M. Emerton. On the interpolation of systems of eigenvalues attached to automorphic Hecke eigenforms. *Invent. Math.*, 164(1):1–84, 2006.
- [25] J.-M. Fontaine. Représentations p -adiques semi-stables. *Astérisque*, (223):113–184, 1994. With an appendix by Pierre Colmez, Périodes p -adiques (Bures-sur-Yvette, 1988).

- [26] M. Harris and R. Taylor. *The geometry and cohomology of some simple Shimura varieties*, volume 151 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2001. With an appendix by Vladimir G. Berkovich.
- [27] L. Herr. Sur la cohomologie galoisienne des corps p -adiques. *Bull. Soc. Math. France*, 126(4):563–600, 1998.
- [28] K. S. Kedlaya. A p -adic local monodromy theorem. *Ann. of Math. (2)*, 160(1):93–184, 2004.
- [29] K. S. Kedlaya and R. Liu. On families of φ, Γ -modules. *Algebra Number Theory*, 4(7):943–967, 2010.
- [30] K. S. Kedlaya, J. Pottharst, and L. Xiao. Cohomology of arithmetic families of (φ, Γ) -modules. *J. Amer. Math. Soc.*, 27(4):1043–1115, 2014.
- [31] M. Kisin. Overconvergent modular forms and the Fontaine–Mazur conjecture. *Invent. Math.*, 153(2):373–454, 2003.
- [32] J.-P. Labesse. Changement de base CM et séries discrètes. In *On the stabilization of the trace formula*, volume 1 of *Stab. Trace Formula Shimura Var. Arith. Appl.*, pages 429–470. Int. Press, Somerville, MA, 2011.
- [33] R. Liu. Cohomology and duality for (ϕ, Γ) -modules over the Robba ring. *Int. Math. Res. Not. IMRN*, (3):Art. ID rnm150, 32, 2008.
- [34] R. Liu. Triangulation of refined families. *Comment. Math. Helv.*, 90(4):831–904, 2015.
- [35] B. Mazur. An introduction to the deformation theory of Galois representations. In *Modular forms and Fermat’s last theorem (Boston, MA, 1995)*, pages 243–311. Springer, New York, 1997.
- [36] B. Mazur. The theme of p -adic variation. In *Mathematics: frontiers and perspectives*, pages 433–459. Amer. Math. Soc., Providence, RI, 2000.
- [37] A. Mínguez. Unramified representations of unitary groups. In *On the stabilization of the trace formula*, volume 1 of *Stab. Trace Formula Shimura Var. Arith. Appl.*, pages 389–410. Int. Press, Somerville, MA, 2011.
- [38] J. Pottharst. Analytic families of finite-slope Selmer groups. *Algebra Number Theory*, 7(7):1571–1612, 2013.
- [39] R. Rouquier. Caractérisation des caractères et pseudo-caractères. *J. Algebra*, 180(2):571–586, 1996.
- [40] S. W. Shin. Galois representations arising from some compact Shimura varieties. *Ann. of Math. (2)*, 173(3):1645–1741, 2011.
- [41] F. Tan. *Families of p -adic Galois Representations*. ProQuest LLC, Ann Arbor, MI, 2011. Thesis (Ph.D.)—Massachusetts Institute of Technology.
- [42] J. T. Tate. Number theoretic background. In *Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2*, Proc. Sympos. Pure Math., XXXIII, pages 3–26. Amer. Math. Soc., Providence, R.I., 1979.
- [43] R. Taylor. Galois representations associated to Siegel modular forms of low weight. *Duke Math. J.*, 63(2):281–332, 1991.
- [44] J. Thorne. On the automorphy of l -adic Galois representations with small residual image. *J. Inst. Math. Jussieu*, 11(4):855–920, 2012. With an appendix by Robert Guralnick, Florian Herzig, Richard Taylor and Thorne.

JOHN BERGDALL, DEPARTMENT OF MATHEMATICS AND STATISTICS, BOSTON UNIVERSITY, 111 CUMMINGTON MALL, BOSTON, MA 02215, USA

E-mail address: bergdall@math.bu.edu

URL: <http://math.bu.edu/people/bergdall>