

Bryn Mawr College

Scholarship, Research, and Creative Work at Bryn Mawr College

Biology Faculty Research and Scholarship

Biology

2021

Studies of insect temporal trends must account for the complex sampling histories inherent to many long-term monitoring efforts

Ellen A.R. Welte

Senckenberg Research Institute and Natural History Museum Frankfurt

Anthony Joern

Kansas State University

Aaron M. Ellison

Harvard University

David C. Lightfoot

University of New Mexico

Sydne Record

Bryn Mawr College, srecord@brynmawr.edu

See next page for additional authors

Follow this and additional works at: https://repository.brynmawr.edu/bio_pubs



Part of the [Biology Commons](#), and the [Ecology and Evolutionary Biology Commons](#)

[Let us know how access to this document benefits you.](#)

Custom Citation

Welte, E.A.R., Joern, A., Ellison, A.M. et al. "Studies of insect temporal trends must account for the complex sampling histories inherent to many long-term monitoring efforts." *Nat Ecol* 5: 589–591.

This paper is posted at Scholarship, Research, and Creative Work at Bryn Mawr College.
https://repository.brynmawr.edu/bio_pubs/41

For more information, please contact repository@brynmawr.edu.

Authors

Ellen A.R. Welti, Anthony Joern, Aaron M. Ellison, David C. Lightfoot, Sydne Record, Nicholas Rodenhouse, Emily H. Stanley, and Michael Kaspari

1 **Matters Arising**

2 **Response to Crossley et al. 2020**

3

4 **Title**

5 Studies of insect temporal trends must account for the complex sampling histories inherent to
6 many long-term monitoring efforts

7

8 **Authors**

9 Ellen A. R. Welte^{1*}, Anthony Joern², Aaron M. Ellison³, David C. Lightfoot⁴, Sydne Record⁵,
10 Nicholas Rodenhouse⁶, Emily H. Stanley⁷, Michael Kaspari⁸

11

12 **Author Affiliations**

13 ¹ Senckenberg Research Institute and Natural History Museum Frankfurt, Gelnhausen, Germany

14 ² Division of Biology, Kansas State University, Manhattan, KS, USA

15 ³ Harvard Forest, Harvard University, Petersham, MA, USA

16 ⁴ Museum of Southwestern Biology, Biology Department, University of New Mexico,
17 Albuquerque, NM, USA

18 ⁵ Department of Biology, Bryn Mawr College, Bryn Mawr, PA, USA

19 ⁶ Biological Sciences, Wellesley College, Wellesley, MA, USA

20 ⁷ Center for Limnology, University of Wisconsin-Madison, Madison, WI, USA

21 ⁸ Geographical Ecology Group, Department of Biology, University of Oklahoma, Norman, OK,
22 USA

23 *corresponding author; email: ellen.welte@senckenberg.de

24

25 **Matters Arising**

26 Crossley et al. (2020)¹ examine patterns of change in insect abundance and diversity across US
27 Long-Term Ecological Research (LTER) sites, concluding “a lack of overall increase or
28 decline”. This is notable if true, given mixed conclusions in the literature regarding the nature
29 and ubiquity of insect declines across regions and insect taxonomic groups²⁻⁶. The data analyzed,
30 downloaded from and collected by US LTER sites, represent unique time series of arthropod
31 abundances. These long-term datasets often provide critical insights, capturing both steady
32 changes and responses to sudden unpredictable events. However, a number of the included
33 datasets are not suitable for estimating long-term observational trends because they come from
34 experiments or have methodological inconsistencies. Additionally, long-term ecological datasets
35 are rarely uniform in sampling effort across their full duration as a result of the changing goals
36 and abilities of a research site to collect data⁷. We suggest that Crossley et al.’s results rely upon
37 a key, but flawed, assumption, that sampling was collected “in a consistent way over time within
38 each dataset”. We document problems with data use prior to statistical analyses from eight LTER
39 sites due to datasets not being suitable for long-term trend estimation and not accounting for
40 sampling variation, using the Konza Prairie (KNZ) grasshopper dataset (CGR022) as an
41 example.

42

43 *Unsuitable datasets to estimate long-term observational trends*

44 Several of the LTER datasets included in Crossley et al. (2020) either document experiments
45 which have confounding treatment effects or they are too variable in sampling methods to allow
46 for comparison of samples across time. Additionally, in one case, Lepidopteran outbreak
47 dynamics with long intervals (10-13 years) at Hubbard Brook limit power to detect meaningful

48 trends without extremely long-term data⁸. Datasets from Cedar Creek include arthropods
49 collected in plots with nitrogen addition, herbivore exclosures, and manipulated plant diversity.
50 All three of the datasets from Harvard Forest included in Crossley et al.'s analysis have large
51 methodological inconsistencies over time and one dataset documents ants collected in a canopy
52 manipulation experiment, including one treatment where trees were girdled to simulate hemlock
53 woolly adelgid (*Adelges tsugae*) infestation of the hemlock trees years prior to the arrival of the
54 invasive insect to the area. One dataset from North Temperate Lakes documents the responses of
55 two crayfish species in a lake where one species was being experimentally removed. With a few
56 exceptions for partial components of these datasets (e.g. control plots in the arce153 Cedar Creek
57 dataset), these data are inappropriate for estimation of long-term observational species trends.


58

59 ***Not accounting for sampling variation: Konza grasshoppers as a case in point***

60 The KNZ CGR022 dataset documents grasshopper species abundances on 15 KNZ watersheds,
61 and spans 1982-present (up to 2015 included in Crossley et al. 2020). Crossley et al. analyze
62 time series of individual species from each dataset (the number of "Time trends" in their Table
63 1). However, regardless of variant sampling effort, they regularly sum all individuals within
64 LTER datasets to yield a single value of abundance for a given species and year. This is the case
65 for KNZ grasshoppers, and most other included datasets (number of "Sites" in their Table 1).
66 Importantly, sampling effort at KNZ and other LTER sites was not constant. At KNZ, variation
67 occurred in the number of samples per watershed and the number of watersheds in which
68 grasshoppers were collected per year (Fig. 1). Most notably, 6 bison-grazed watersheds were
69 added to KNZ sampling in 2002. Changes in sample numbers over time are documented in the

70 online metadata (<http://lter.konza.ksu.edu/content/cgr02-sweep-sampling-grasshoppers-konza-prairie-lter-watersheds>).

72



73

74 **Figure 1. The complex history of sampling of the KNZ grasshopper dataset.** The KNZ

75 grasshopper dataset (CGR022) exhibits high variance both in number of watersheds sampled per
 76 year (number of bars per year) and number of samples collected within each watershed each year
 77 (depicted in color). Other complexities include the tragic loss of four years (1992-1995) of
 78 sampling due to a freezer crash, changes in sampling month, changes in watershed burn
 79 frequencies, and the reintroduction of bison in the 1990s to six of the later-sampled watersheds.

80

81 Accounting for sampling effort and data structure matters (*see also* Supplementary
 82 Information: Fig. S1). At KNZ, bison-grazed watersheds support higher grasshopper abundances
 83 and species richness^{9,10}. In a recent analysis using the CGR022 dataset, to account for this
 84 change in sampling effort, data were combined only from watersheds collected in the same years
 85 (e.g. by splitting samples from grazed watersheds into a separate time series) and abundances

86 within each watershed and year were divided by the number of samples. Analysis of the data
87 structured in this way showed a >2% annual decline in grasshopper abundance, with only one
88 common species increasing¹¹. Crossley et al., in contrast report most grasshopper species
89 increased in abundance from 1982-2015. The authors of Crossley et al. (2020) note the
90 discrepancy with both this study¹¹ and another³, and suggest it is “driven by falling numbers of
91 just two once-dominant species... whereas many other formerly rare species have become more
92 abundant and both evenness and species richness have increased”. However, we believe the
93 discrepancy arises because Crossley et al. did not account for variable sampling effort, including
94 KNZ’s incorporation of additional, more diverse grazed habitats midway in the time series.
95 Similar errors, where data structure was not accounted for, are evident in 17 of the 19 datasets
96 which we examined and were included in Crossley et al. (2020)’s results.

97

98 ***Conclusion***

99 We have thus far been able to confirm issues with data from 8 of the 13 LTER sites (comprising
100 60% of Table 1’s “Time trends”) included in Crossley et al. (2020). We note that this is not a
101 comprehensive assessment, as we have only included errors from datasets of which either we
102 ourselves are the PIs or we have been able to confirm with the corresponding LTER PIs and
103 information managers. The eight sites are: Baltimore, Cedar Creek, Central Arizona-Phoenix,
104 Harvard Forest, Hubbard Brook, Konza Prairie, North Temperate Lakes, and Sevilleta. We
105 provide details on dataset unsuitability, mistakes in not accounting for sampling effort, and
106 several coding errors in the Supplementary Information.

107

108 Given these mistakes, we urge skepticism regarding Crossley et al. (2020)'s general
109 conclusion of no net decline in insect abundances at US LTER sites in recent decades. Although
110 their goal is laudable, both the use of unsuitable datasets and not taking sampling effort into
111 account generate erroneous estimates of population change. Recently, a study reporting
112 widespread collapse of rainforest insect populations at the LTER site Luquillo necessitated a
113 similar correction⁵. We echo those authors, when they suggest that scientists can avoid errors by
114 reading corresponding metadata. Contacting in advance (or even including as authors) the data
115 providers/field biologists are additionally good practices that ensure appropriate use of the data.
116 Like the ecology they document, it is important to take into account that long-term monitoring
117 efforts by LTERs and similar institutions are themselves complex and full of history.

118

119 **Author Contributions**

120 E.A.R.W., S. R., A.J., and M.K. conceived the idea for the paper. E.A.R.W. wrote the first draft.
121 A.M.E., D.L., S.R., N.R., and E.S. identified further errors in the Crossley et al. online data. All
122 authors significantly contributed to revisions.

123

124 **Acknowledgements**

125 Shannon LaDeau, Stevan Earl, Susan Barrott, Elizabeth Borer, and Steve Pennings aided in
126 identifying errors in Crossley et al.'s online data. Karl Roeder provided comments on an early
127 draft of this manuscript. Jeff Taylor aided in identifying changes in Konza watershed names. We
128 thank all LTER information managers and PIs who help keep online metadata updated. We thank
129 Pam Montz, John Haarstad, Amanda Kuhl, Matthew Ayres, Richard Holmes, and the numerous

130 others who did the hard work to generate these long-term datasets. NSF DEB-1556280 to MK
131 and Konza Prairie LTER NSF DEB-1440484 supported this work.

132

133 **Competing Interests**

134 The authors declare no competing interests.

135

136 **Data Availability**

137 KNZ grasshopper abundance data are available from the Long-Term Ecological Research Data
138 Portal (<https://doi.org/10.6073/pasta/7b2259dcb0e499447e0e11dfb562dc2f>). Citations for the
139 additionally described LTER datasets are provided in the Supplementary Information.

140

141 **References**

- 142 1. Crossley, M. S. *et al.* No net insect abundance and diversity declines across US Long Term
143 Ecological Research sites. *Nature Ecology & Evolution* (2020) doi:10.1038/s41559-020-1269-
144 4.
- 145 2. Hallmann, C. A. *et al.* More than 75 percent decline over 27 years in total flying insect
146 biomass in protected areas. *PLoS One* **12**, e0185809 (2017).
- 147 3. van Klink, R. *et al.* Meta-analysis reveals declines in terrestrial but increases in freshwater
148 insect abundances. *Science* **368**, 417–420 (2020).
- 149 4. Lister, B. C. & Garcia, A. Climate-driven declines in arthropod abundance restructure a
150 rainforest food web. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E10397–E10406 (2018).

- 151 5. Willig, M. *et al.* Populations are not declining and food webs are not collapsing at the
152 Luquillo Experimental Forest. *Proceedings of the National Academy of Sciences* **116**, 12143–
153 12144 (2019).
- 154 6. Pilotto, F. *et al.* Meta-analysis of multidecadal biodiversity trends in Europe. *Nature*
155 *Communications* **11**, 3486 (2020).
- 156 7. Didham, R. K. *et al.* Interpreting insect declines: seven challenges and a way forward. *Insect*
157 *Conservation and Diversity* **13**, 103–114 (2020).
- 158 8. Martinat, P. J. & Allen, D. C. Saddled prominent outbreaks in North America. *Northern*
159 *Journal of Applied Forestry* **5**, 88–91 (1988).
- 160 9. Welti, E. A. R. *et al.* Fire, grazing and climate shape plant-grasshopper interactions in a
161 tallgrass prairie. *Funct. Ecol.* **33**, 735–745 (2019).
- 162 10. Bruckerhoff, L. A. *et al.* Harmony on the prairie? Grassland plant and animal community
163 responses to variation in climate across land-use gradients. *Ecology* **101**, e02986 (2020).
- 164 11. Welti, E. A. R., Roeder, K. A., de Beurs, K. M., Joern, A. & Kaspari, M. Nutrient
165 dilution and climate cycles underlie declines in a dominant insect herbivore. *Proc. Natl. Acad.*
166 *Sci. U. S. A.* **117**, 7271–7275 (2020).
- 167