

Bryn Mawr College

Scholarship, Research, and Creative Work at Bryn Mawr College

Psychology Faculty Research and Scholarship

Psychology

2016

Differential effects of mineralocorticoid and angiotensin II on incentive and mesolimbic activity

Laura A. Grafe

Bryn Mawr College, lgrafe@brynmawr.edu

Loretta M. Flanagan-Cato

Follow this and additional works at: https://repository.brynmawr.edu/psych_pubs



Part of the [Psychology Commons](#)

[Let us know how access to this document benefits you.](#)

Citation

Grafe, Laura A. and Loretta M. Flanagan-Cato. 2016. "Differential effects of mineralocorticoid and angiotensin II on incentive and mesolimbic activity." *Hormones and Behavior* 79: 28–36.

This paper is posted at Scholarship, Research, and Creative Work at Bryn Mawr College.
https://repository.brynmawr.edu/psych_pubs/71

For more information, please contact repository@brynmawr.edu.

1
2 **DIFFERENTIAL EFFECTS OF MINERALOCORTICOID**
3 **AND ANGIOTENSIN II ON INCENTIVE AND MESOLIMBIC ACTIVITY**

4
5 Laura A. Grafe^a and Loretta M. Flanagan-Cato^{a,b,c}

6
7 Neuroscience Graduate Group^a, Department of Psychology^b, and the Mahoney Institute of
8 Neurological Sciences^c, University of Pennsylvania,

9 Philadelphia Pennsylvania, USA

10
11 Abbreviated title: Aldo and AngII-induced motivation for sodium

12
13
14 Corresponding Author:

15 L.M. Flanagan-Cato

16 flanagan@psych.upenn.edu

17 3720 Walnut Street, Philadelphia, PA, 19104

18
19 Laura A. Grafe co-designed the research, executed the experiments, analyzed the data, and co-
20 wrote the paper. Loretta M. Flanagan-Cato co-designed research, analyzed the data, and co-
21 wrote the paper.

22

23 ABSTRACT

24 The controls of thirst and sodium appetite are mediated in part by the hormones
25 aldosterone and angiotensin II (AngII). The present study examined the behavioral and
26 neural mechanisms of altered effort-value in animals treated with systemic
27 mineralocorticoids, intracerebroventricular AngII, or both. First, rats treated with
28 mineralocorticoid and AngII were tested in the progressive ratio operant task. The
29 willingness to work for sodium versus water depended on hormonal treatment. In
30 particular, rats treated with both mineralocorticoid and AngII preferentially worked for
31 access to sodium versus water compared with rats given only one of these hormones.
32 Second, components of the mesolimbic dopamine pathway were examined for
33 modulation by mineralocorticoids and AngII. Based on cFos immunohistochemistry,
34 AngII treatment activated neurons in the ventral tegmental area and nucleus accumbens,
35 with no enhancement by mineralocorticoid pretreatment. In contrast, western blot
36 analysis revealed that combined hormone treatment increased levels of phospho-tyrosine
37 hydroxylase in the ventral tegmental area. Thus, mineralocorticoid and AngII treatments
38 differentially engaged the mesolimbic pathway based on tyrosine hydroxylase levels
39 versus cFos activation.

40

41 Key words: Aldosterone; Angiotensin; Dopamine; Motivation; Nucleus Accumbens; Sodium
42 Appetite; Thirst; Ventral Tegmental Area

43

44

45 **Introduction**

46

47 Fluid depletion can be life threatening, and animals must carefully titrate their
48 intake of water and sodium to restore and maintain osmotic and volemic balance.
49 Sodium replacement requires the goal-directed behavior known as sodium appetite
50 (Andersson, 1977), a behavior that can be prompted by mineralocorticoids, such as
51 aldosterone, and angiotensin II (AngII) (Johnson and Thunhorst, 1997). AngII acts in the
52 brain to elicit water intake and sodium intake (Epstein et al., 1969). During fluid depletion,
53 suppression of either central AngII or aldosterone action does not eliminate sodium
54 appetite, but blocking the central actions of both hormones abolishes the behavior (Buggy
55 and Jonklaas, 1984; Sakai et al., 1986). Conversely, when both aldosterone and AngII
56 are given exogenously, sodium appetite is potentiated (Fluharty and Epstein, 1983). The
57 behavioral and neural basis for the combined effect of aldosterone and AngII on sodium
58 appetite remains undefined.

59 The behavioral effects aldosterone and AngII to promote sodium ingestion may
60 involve parallel behavioral mechanisms. For example, sodium ingestion could be
61 enhanced by a change in the hedonic strength of the sodium tastant. Indeed, in rats
62 placed on a sodium deficient diet, which increases aldosterone and AngII levels, sodium
63 ingestion is preferred to moderately reinforcing brain stimulation, which suggests sodium
64 appetite involves the modulation of the pleasurable properties of sodium intake (Conover
65 et al., 1994). In parallel to altered taste value, sodium appetite may involve a recalibration
66 of incentive-based effort. In this regard, rats treated with both aldosterone and AngII run
67 faster on a runway to gain access to sodium, compared with rats treated with either

68 hormone alone (Zhang et al., 1984), which suggests that the combined hormone
69 treatment increases the incentive value of sodium. The progressive ratio task is a
70 quantitative assay for incentive-based effort, but the effects of combined
71 mineralocorticoids and AngII on this behavioral test have not been reported.

72 The mesolimbic dopamine system has been widely implicated in effort-related
73 behaviors (Barbano and Cador, 2006; Floresco et al., 2008; Kelley et al., 2005; Phillips
74 et al., 2007; Salamone et al., 2009). Dopamine neurons in the ventral tegmental area
75 (VTA) project to the accumbens, which in turn projects to brain regions such as the ventral
76 pallidum to generate goal directed movement (Carelli, 2002). Previous work has
77 implicated this brain system in sodium appetite. For example, the accumbens receives
78 multisynaptic input from aldosterone-sensing and sodium-sensing neurons in the
79 hindbrain (Miller and Loewy, 2014; Shekhtman et al., 2007). Sodium depletion alters the
80 level of dopamine transporters and opioid peptides in the accumbens (Grondin et al.,
81 2011; Lucas et al., 2003; Roitman et al., 1999). In addition, sodium depletion modifies
82 the dendritic arbor of ventral striatum neurons (Roitman et al., 2002). Although these
83 studies have suggested a role for mesolimbic activity in sodium appetite, the separate
84 effects of aldosterone and AngII on mesolimbic activation have not been studied.

85 The present studies tested the overall hypothesis that mineralocorticoids and AngII
86 recalibrate the willingness to work for sodium versus water. In particular, the willingness
87 to work was measured with the progressive ratio task. In addition, the activity of the
88 mesolimbic dopamine pathway was assessed with immunohistochemistry for cFos and
89 western blot analysis for tyrosine hydroxylase. Although it is recognized that sodium
90 depletion is a complex physiological state that is imperfectly mimicked with

91 mineralocorticoid and central AngII treatments, this preparation has yielded useful
92 insights into the neuroendocrine actions that influence motivated behavior.

93

94 **Materials and Methods**

95 *Animals*

96 Adult male Sprague-Dawley rats (weight between 225 and 250 g) were obtained
97 from Charles River Laboratories (Wilmington, MA, USA). Rats were pair-housed in plastic
98 tubs with standard bedding and with food and water available *ad libitum*, except during
99 experimental procedures. The temperature in the colony was maintained at 22°C with a
100 12:12 h reversed light/dark cycle. Behavioral testing, described below, was conducted
101 during the lights-out phase. Animals were allowed at least one week to acclimate to the
102 colony before any procedures were performed. The Institutional Animal Care and Use
103 Committee of the University of Pennsylvania approved all procedures with animals.

104

105 *Surgery*

106 Surgeries were performed in aseptic conditions. Animals were anesthetized with
107 inducted and maintained with isoflurane anesthesia for stereotaxic surgery. A 26-gauge
108 guide cannula (Plastics One, Roanoke, VA, USA) was aimed at the lateral ventricle using
109 these coordinates: 0.48 mm caudal to bregma, 1.6 mm from mid-line and 4.2 mm ventral
110 to dura mater. The cannulae were fixed in place with dental cement and bone screws.
111 Upon completion of surgery procedures, animals were injected with yohimbine (0.11

112 mg/kg, ip, Ben Venue Laboratories Bedford, OH), and upon awakening animals were
113 returned to the housing facility and singly housed. The animals were allowed at least five
114 days to recover before verification procedures were performed.

115 Prior to undergoing experimental treatments, animals were tested for correct
116 lateral ventricle cannula placement and patency. They were given an *icv* injection of 20.0
117 ng of AngII diluted in artificial cerebrospinal fluid (aCSF) via a Hamilton syringe connected
118 with PE-10 tubing to an injector that terminated 1 mm beyond the guide cannula. Animals
119 were excluded from the experiment if they failed to demonstrate a drinking response in
120 less than 30 seconds, consuming at least 3 ml of water, in two separate AngII challenges.
121 Experiments began three days after these *icv* test injections.

122

123 *Experimental Design*

124 In all experiments, animals were assigned to one of four treatments in a 2 x 2
125 design, with a crossover in behavioral experiments. Animals were first pre-treated twice
126 daily (10 hours apart) for three days with a subcutaneous injection of sesame oil or an
127 aldosterone analog, deoxycorticosterone acetate (DOC; 0.25 mg/0.2 ml sesame oil;
128 Sigma, St Louis, MO). DOC penetrates the blood brain barrier more easily than
129 aldosterone due to its low capacity for hydrogen bond formation (Kraulis et al., 1975).
130 Animals then were injected *icv* with either artificial cerebrospinal fluid (aCSF; R&D
131 Systems, Minneapolis, MN) or 20.0 ng AngII in a volume of 2.0 ul (Bachem, King of
132 Prussia, PA). The treatment groups will be referred to as follows: Veh/Veh, DOC/Veh,
133 Veh/AngII, DOC/AngII. Using this experimental design and identical doses, the

134 DOC/AngII treatment has been shown to elicit a greater than additive effect on sodium
135 intake, but not water intake (Grafe et al., 2014).

136

137 *Experiment 1. Progressive Ratio Task*

138 Rats were acclimated to wire mesh cages for one hour with two 25-ml bottles
139 containing tap water and 3% saline, each marked with 0.2 ml graduations. These bottles
140 were then removed, and rats were water restricted for 23 hours per day for the next six
141 days. During these six days, rats underwent operant lever pressing training in
142 conditioning boxes for 30 minutes per day (Med Associates; MDPC IV Software, St.
143 Albans, Vermont). The conditioning boxes contained levers for both water (right lever)
144 and 3% saline (left lever), both simultaneously present. A lever press lowered a syringe
145 pump, which delivered a 0.1 ml drop of the appropriate liquid into a cup available to the rat. The
146 saline and water each had their own syringe pump and their own cup. During the first two
147 training days, to facilitate learning, an aliquot was dispensed every 300 sec that elapsed
148 without bar pressing. In addition, animals could earn a 0.1 ml of water or saline for each
149 bar press, depending on which of the two levers was pressed. During the subsequent
150 two training days, the animals earned a 0.1 ml water or saline for each lever press,
151 followed by two training days during which three lever presses were required for each
152 aliquot of water or saline. Animals were considered to have learned the lever-fluid
153 contingencies when they had made at least 10 lever presses for water during the 30-min
154 session. Once this occurred, rats were given ad libitum access to water again. Rats were
155 then assigned to treatment groups, as described above, and given no further operant

156 training while they received their three days of pretreatment injections (vehicle or DOC).
157 After pretreatment was complete, 24 hours after the last DOC treatment, rats were
158 administered their assigned *icv* injection (vehicle or 20 ng AngII), and immediately given
159 a test with a progressive ratio (PR) schedule. Thus, animals were water replete at the
160 beginning of the PR test. The response requirement of the PR schedule increased
161 progressively for both saline and water, as previously described (Davis et al., 2011). The
162 breakpoint for each animal was defined as the final reinforced bar pressing set that
163 preceded a 10-minute period without earning a reinforcement, with a two-hour limit total.
164 Food was not available during this task.

165

166 *Experiment 2: Hormone-Induced cFos expression*

167 To observe brain activation after DOC and AngII treatments, rats were assigned
168 to treatment groups, as explained above. Sixty minutes after the last *icv* injection, each
169 rat was anesthetized with 50 mg/kg ketamine and 20 mg/kg xylazine, intraperitoneally.
170 As discussed below, there were group differences in the effort for sodium versus water
171 during the first few minutes of the progressive ratio task, making 60 minutes post-
172 treatment a reasonable time to expect differences in cFos levels. Rats were perfused
173 transcardially with 100 mL of heparinized saline followed by 200 ml 4% paraformaldehyde
174 (Electron Microscopy Sciences, Fort Washington, PA). The brains were isolated, post-
175 fixed in paraformaldehyde overnight at 4°C, and then submerged in 20% sucrose in 0.1
176 M phosphate buffer for three days. Coronal sections were cut on a freezing microtome
177 into three serial sets of 40-um-thick sections. These sections encompassed the VTA and

178 the shell and core of the accumbens. One set of sections from each animal underwent
179 immunohistochemical staining and analysis; the other two sets of sections were
180 preserved in cryoprotectant as reserve material.

181 Sections were washed in Tris-buffered saline (TBS; pH 7.4) and then incubated
182 with a cFos antibody (1:500, sc-52, rabbit; Santa Cruz Biotechnology, Santa Cruz, CA) in
183 TBS with 0.2% TritonX-100 and 3% normal donkey serum (Jackson ImmunoResearch;
184 West Grove, PA) overnight at 4°C. After several washes, sections were incubated with a
185 Biotin-SP-conjugated AffiniPure Donkey Anti-rabbit IgG (1:100, Jackson
186 ImmunoResearch) in TBS with 0.2% TritonX-100 and 3% normal donkey serum for 2 hours
187 at room temperature. After several washes, sections were incubated with the Vectastain
188 ABC kit (Vector Laboratories, Burlingame, CA) for one hour. This was followed by another
189 set of washes before staining with 3'3'-diaminobenzidine (Sigma-Aldrich) for 10 minutes.
190 After a final set of washes, sections were mounted on slides, air-dried, and cover slipped
191 with DPX mounting media (Electron Microscopy Sciences: Fort Washington, PA).

192 Photomicrographs were acquired with a digital camera (Diagnostic Instruments,
193 Sterling Heights, MI, model RTKE), maintaining the same microscope and camera
194 settings to ensure the same level of light and exposure for all images. Background was
195 subtracted from images using Photoshop, and images were set to the same threshold
196 level (200). Images were further analyzed in NIH Image J using standardized boxes for
197 each brain region (based on Paxinos & Watson Rat Brain Atlas, shown in **Figure 1A**),
198 using the analyze particles function, which counts objects within specific size and shape
199 parameters. Pixel size minimum was 15 and circularity was set to 0.35.

200

201 *Experiment 3: Hormone Regulation of Tyrosine Hydroxylase*

202 Animals were assigned to four treatment groups, as described above. Five
203 minutes after the last injections, animals were rapidly decapitated and brains were flash
204 frozen in hexane over dry ice. The five-minute time point was selected based on its
205 correlation with a significant increase in lever presses for sodium in the progressive ratio
206 experiment. Using a cryostat, 1-mm punches of the brain regions of interest (VTA,
207 accumbens shell and core, shown in **Figure 1B**) were collected from 300- μ m sections.
208 Punches were immersed in lysis buffer containing 25 mM Tris-HCl (pH 8), protease
209 inhibitors (pepstatin, leupeptin, aprotinin), and phosphatase inhibitors (sodium
210 pyrophosphate, sodium fluoride, sodium molybdate, phenylarsin oxide, and sodium
211 orthovanadate). The VTA and accumbens (shell and core) punches were immersed in
212 50 and 100 μ L of lysis buffer, respectively. Brain punches were sonicated for three
213 seconds followed by centrifugation at 14,000 rpm in 4°C for 15 minutes. Supernatant was
214 collected and a Bicinchoninic acid (BCA) protein assay was performed on five microliters
215 of each sample. Based on the protein levels detected by the BCA assay, appropriate
216 amounts of sample and sample buffer were loaded into wells of a 10% sodium dodecyl
217 sulfate polyacrylamide gel. Western blots for phospho- and total TH were performed
218 using the LiCor Odyssey System. The following antibodies were used: monoclonal anti-
219 tyrosine hydroxylase antibody T2928 at 1:8000 (Sigma Aldrich); tyrosine hydroxylase
220 pS31 rabbit polyclonal antibody #36-9900 at 1:400 (Life Technologies, Carlsbad, CA);
221 IRDye 800CW Goat anti-Mouse IgG (H + L) 926-32210 at 1:2000 (Li-Cor Biosciences,
222 Lincoln, NE); IRDye 680LT Goat anti-Rabbit IgG (H + L) 926-68021 at 1:4000 (Li-Cor

223 Biosciences). Phosphorylation sites on tyrosine hydroxylase generally increase
224 catecholamine production. Serine 31, the residue attended to in the present study, is
225 phosphorylated by depolarization and extracellular receptor-activated kinases 1 and 2
226 (Tekin et al., 2014).

227

228 *Statistical Analysis*

229 Data are presented as the mean \pm the standard deviation of the mean. For all
230 experiments, comparisons were made between treatment variables with a two-way
231 ANOVA. When warranted, Bonferonni-corrected t-tests were performed. Effect sizes
232 were calculated using η^2 and Cohen's d. All hypothesis tests used $\alpha=0.05$ as the criterion
233 level of significance. Statistical analyses were conducted using Prism 2.0 software (La
234 Jolla, CA).

235

236 **Results**

237

238 The hormone regimen used in the present study was previously demonstrated to
239 increase sodium ingestion in the Veh/AngII and DOC/AngII groups and to significantly
240 water intake in the Veh/AngII condition (Grafe et al., 2014). Here, the progressive ratio
241 operant task was used to compare the effort rats are willing to exert for sodium ingestion
242 after these hormone treatments. Rats were pretreated with oil or DOC, followed by *icv*
243 treatments of vehicle or AngII and allowed access to two levers: one for access to 3%
244 saline and the other for access to water (0.1 ml per reinforcement). **Figure 2A** illustrates

245 the cumulative number of lever presses across the duration of testing for 3% saline and
246 water. DOC/AngII treated rats bar-pressed for a longer duration for 3% saline compared
247 with the other treatment groups.

248 We conducted separate two-way ANOVA tests to examine effects of these
249 hormones on bar presses for sodium and water. The two-way ANOVA supported a main
250 effect for AngII ($F(1,29) = 13.9$, $p = 0.0008$, $\eta^2 = 0.31$) but not DOC ($F(1,29) = 0.1$, $p =$
251 0.752 , $\eta^2 > 0.01$) on 3% saline lever presses. Bonferroni-corrected post-hoc t tests
252 revealed that the DOC/AngII group pressed significantly more on the saline lever
253 compared with the Veh/Veh and DOC/Veh groups (Veh/Veh: 10.9 presses (SD2.4),
254 DOC/Veh: 6.4 presses (SD1.7), Veh/AngII: 19.1 presses (SD6.4), DOC/AngII: 26.0
255 presses (SD2.8); $p = 0.001$, $p = 0.0001$; $d = 1.50$). A two-way ANOVA established a main
256 effect for AngII ($F(1,29) = 5.7$, $p = 0.023$, $\eta^2 = 0.13$) but not DOC ($F(1,29) = 1.7$, $p = 0.203$,
257 $\eta^2 = 0.04$) on water lever presses, and a significant interaction between the two hormones
258 ($F(1,29) = 8.4$, $p = 0.007$, $\eta^2 = 0.19$). Post-hoc analysis revealed that rats treated with
259 Veh/AngII pressed more for water than any other treatment group (Veh/Veh: 16.9 presses
260 (SD5.2), DOC/Veh: 29.5 presses (SD4.9), Veh/AngII: 58.9 presses (SD10.7),
261 DOC/AngII: 25.5 presses (SD8.9); $p = 0.003$, $p = 0.020$, $p = 0.030$; $d = 1.47$).

262 With a progressive ratio schedule, the response requirement to attain rewards
263 increases according to the same rule throughout the test session until the rat stops
264 responding (Sclafani and Ackroff, 2003). The highest set of bar presses completed is
265 referred to as the break point and provides a measure of incentive. In the present
266 experiment, rats had simultaneous access to work for sodium and water. Treatments for
267 Veh/Veh, DOC/Veh and Veh/AngII elicited more work for water compared with sodium,

268 whereas the DOC/AngII-treated animals exhibited the opposite pattern. A two-way
269 ANOVA established an interaction between AngII and DOC ($F(1,29) = 4.6, p = 0.046$;
270 $F(1,29) = 8.6, p = 0.006; \eta^2 = 0.18$). DOC/AngII treated rats had significantly higher ratio
271 of the breakpoint for sodium versus breakpoint for water than all other treatment groups.

272

273 *Experiment 2: Hormone-Induced cFos expression*

274 The observed unique effect of DOC/AngII treatment on willingness to work for
275 sodium versus water suggested an underlying difference in mesolimbic activity. As
276 shown in **Figure 3**, the number of cFos labeled cells was quantified in the VTA and the
277 accumbens after Veh/Veh, DOC/Veh, Veh/AngII, and DOC/AngII treatment. A two-way
278 ANOVA supported a main effect for AngII ($F(1,13) = 39.94, p < 0.0001, \eta^2 = 0.64$), but
279 not for DOC or an interaction between the two hormones in the VTA. Post hoc tests
280 indicated that Veh/AngII and DOC/AngII treatments increased cFos expression compared
281 with Veh/Veh ($p = 0.0001, p = 0.0003$, respectively; $d = 3.77$). In the nucleus accumbens
282 core, a two-way ANOVA revealed a main effect for AngII ($F(1,13) = 12.82, p = 0.0034, \eta^2$
283 $= 0.44$), but no effect for DOC or their interaction on cFos expression. Post-hoc tests
284 indicated that Veh/AngII and DOC/AngII treatments increased cFos levels compared to
285 Veh/Veh ($p = 0.0004, p = 0.0046$, respectively; $d = 2.12$). Lastly, a two-way ANOVA for
286 the nucleus accumbens shell revealed a main effect for AngII ($F(1,13) = 18.6, p = 0.0008,$
287 $\eta^2 = 0.52$), but no effect for DOC or the two-hormone interaction on cFos expression.
288 Post-hoc t tests indicated that each AngII treatment group induced significant cFos
289 expression compared to the Veh/Veh treatment (Veh/AngII $p = 0.0012$; DOC/AngII $p =$

290 0.002; $d = 2.48$). To summarize, AngII treatment activated cFos expression in the VTA
291 and nucleus accumbens, but DOC pretreatment did not enhance this activation.

292

293 *Experiment 3: Hormone Regulation of Tyrosine Hydroxylase*

294 The AngII-induced expression of cFos of the VTA observed in Experiment 2
295 suggested a possible increase in dopamine neurotransmission. The level of tyrosine
296 hydroxylase, the rate-limiting enzyme for dopamine production, and its phosphorylation,
297 were measured by western blot analysis. Rats were pretreated with vehicle or DOC
298 followed by *icv* injections of vehicle or AngII, and tissue punches were collected five
299 minutes after the last injection. In the VTA, where dopaminergic neurons reside, a two-
300 way ANOVA supported a main effect for both DOC and AngII on tyrosine hydroxylase
301 levels in the VTA ($F(1,8) = 16.6$, $p = 0.004$, $\eta^2 = 0.51$; $F(1,8) = 6.0$, $p = 0.04$, $\eta^2 = 0.19$),
302 as illustrated in **Figure 4A**. Post hoc tests indicated each treatment group significantly
303 enhanced tyrosine hydroxylase levels compared with vehicle ($p = 0.003$, $p = 0.04$, $p =$
304 0.02 for DOC/Veh, Veh/AngII, and DOC/AngII, respectively; $d = 1.99$). However, DOC
305 pretreatment did not further enhance the effects of AngII on tyrosine hydroxylase levels.
306 For phospho-tyrosine hydroxylase, the value for each animal, as determined by western
307 blot analysis, was divided by the value for total tyrosine hydroxylase (p-TH/TH). In the
308 VTA, a two-way ANOVA supported a main effect for DOC ($F(1,8) = 10.37$, $p = 0.012$, η^2
309 $= 0.52$), but not AngII treatment, as shown in **Figure 4B**. Post-hoc analysis indicated that
310 in the VTA DOC/AngII groups displayed increased p-TH/TH levels compared with
311 Veh/Veh ($p = 0.02$; $d = 0.78$).

312 In the nucleus accumbens core, neither DOC nor AngII had a significant main
313 effect on tyrosine hydroxylase levels. In the shell of the nucleus accumbens, DOC, but
314 not AngII, had a main effect on levels of tyrosine hydroxylase ($F(1,8) = 9.3$, $p = 0.02$, η^2
315 $= 0.40$). Post-hoc analysis revealed DOC/Veh increased tyrosine hydroxylase levels
316 compared with Veh/Veh ($p = 0.04$; $d = 2.17$). In both the core and shell DOC and AngII
317 treatments significantly increased p-TH/TH compared with vehicle (Core: Veh/Veh: 0.02
318 (SD0.2), DOC/Veh: 0.42 (SD0.2), Veh/AngII: 0.37 (SD0.14), DOC/AngII: 0.38 (SD0.01),
319 $p = 0.011$, $p = 0.020$, $p = 0.018$; $d = 0.72$); (Shell: Veh/Veh: 0.07 (SD0.05), DOC/Veh:
320 0.46 (SD0.02), Veh/AngII: 0.38 (SD0.06), DOC/AngII: 0.39 (SD0.04); $p = 0.0008$, $p =$
321 0.0028 , $p = 0.0022$, respectively; $d = 0.51$). In summary, in both regions of the nucleus
322 accumbens, DOC and AngII pretreatments increased phosphorylation levels of tyrosine
323 hydroxylase, usually without detectably increasing total tyrosine hydroxylase levels.

324

325 **Discussion**

326

327 During sodium deficiency, aldosterone and AngII act in concert to produce an avid
328 consumption of sodium (Fluharty and Sakai, 1995), and combined treatment with these
329 hormones is a useful experimental model for investigating the biological basis of this
330 striking example of motivated behavior. The current experiments tested the overall
331 hypothesis that DOC and AngII treatments uniquely affect the willingness to work for
332 access to sodium, potentially by modulating neural activation and neurochemistry in the
333 mesolimbic dopamine pathway. DOC/AngII-treated rats steadily exerted effort for access
334 to 3% sodium solution while decreasing their effort for access to water compared with the

335 Veh/AngII-treated rats. DOC and AngII had differential effects on cFos activation and
336 phospho-tyrosine hydroxylase levels in the VTA. Additive effects of DOC and AngII were
337 not observed when these treatments were combined, although it is possible that cFos
338 immunohistochemistry and tyrosine hydroxylase immunoblots were limited by a ceiling
339 effect. Instead, the co-treatment of DOC and AngII was associated with a unique
340 combination of enhanced dopamine synthetic capacity and neural activation. **Figure 5**
341 summarizes these results according to hormone treatment, brain region and behavior.

342 Several caveats are well known for the interpretation of cFos expression. First,
343 neuronal activation may occur without cFos induction. Thus, the lack of cFos activation
344 after DOC treatment may not reflect the recent activity of these neurons. In addition, the
345 neural consequences of cFos activation are not fully understood, making cFos merely a
346 marker of recent activity. Thirdly, without knowing the phenotype of the cFos labeled
347 neurons in this study, the increased activity may occur in excitatory or inhibitory neurons,
348 which would have quite different physiological consequences. With those points
349 notwithstanding, cFos expression in the VTA and ventral striatum has been a valuable a
350 proxy for measuring neuronal activation in many studies (Kovács, 2008).

351

352 *Behavioral mechanisms*

353 Healthy rats normally avoid concentrated sodium solutions (Berridge et al., 1984);
354 however when levels of both mineralocorticoids and AngII are elevated, a robust sodium
355 appetite emerges (Fluharty and Epstein, 1983; Shade et al., 2002). A component of
356 sodium appetite is altered gustatory hedonic processing, and subsequent increased

357 pleasure, of passively received oral sodium. In the taste reactivity test, sodium appetite
358 is associated with reduced gaping and increased pre-swallowing behaviors in response
359 to orally presented sodium solutions (Berridge et al., 1984; Clark and Bernstein, 2006).
360 Further evidence that sodium depletion increases the “liking” of concentrated sodium
361 solutions is the depletion-specific response of neurons in the ventral pallidum, a brain
362 region thought to encode pleasure, in response to sodium ingestion (Tindell et al., 2006).
363 The ventral forebrain may contribute to the hormonal modulation of sodium as a tastant
364 (Garcia et al., 2008; Geerling and Loewy, 2006; Lundy, 2008).

365 In addition to enhanced palatability, sodium appetite includes a willingness to work
366 for sodium. For example, rats run faster along a runway to gain access to sodium after
367 co-administration of aldosterone and AngII (Zhang et al., 1984). In the progressive ratio
368 test, the highest set of bar presses completed, referred to as the break point, provides a
369 quantification of goal value (Hodos, 1961). Progressive ratio schedules model foraging
370 in a natural setting; as resources are consumed in a given area, more effort is required to
371 obtain them. This method has been used to quantify the motivation for sodium after body
372 fluid depletion (Clark and Bernstein, 2006; Starr and Rowland, 2006). The present
373 experiment used a two-bottle progressive ratio task to document the effects of
374 mineralocorticoids and AngII on the relative effort-value of 3% sodium solution versus
375 water. This preparation revealed a shift in effort-value, with AngII alone biasing the
376 animals to exert more effort for water than sodium, whereas DOC/AngII treatment drove
377 them to work more for 3% saline than water. Unlike previous studies of sodium appetite
378 that focused on the quantity of sodium ingested, the two-bottle progressive ratio task
379 quantifies the competing drives for sodium and water without changing fluid balance.

380 Differences in effort-value led us to investigate effects of mineralocorticoids and AngII on
381 mesolimbic regions, a brain system that regulates effort-related behaviors (Barbano and
382 Cador, 2006; Floresco et al., 2008; Kelley et al., 2005; Phillips et al., 2007; Salamone et
383 al., 2009).

384 In the present study, no simple correlation exists between the level of cFos
385 activation in a specific brain region and behavior. Rather, similar cFos levels were
386 observed in the VTA and nucleus accumbens when animals would have been working
387 for water versus sodium. One interpretation of this finding is that cFos activation in the
388 mesolimbic dopamine pathway represents a non-specific increase in willingness work.
389 However, it remains uncertain whether the similar numbers of cFos labeled cells were of
390 the same cell type. Furthermore, despite the similarities in cFos levels despite during
391 these different motivated states, it should be noted that regulation of tyrosine hydroxylase
392 varies, suggesting differential dopamine transmission. Thus, further studies are needed
393 to understand the details of motivation-specific regulation of mesolimbic dopamine.

394

395 *Hormone-sensing brain regions*

396 The initial brain targets of mineralocorticoids and AngII are known. Regarding
397 mineralocorticoids, the nucleus of the solitary tract contains a unique population of
398 neurons that co-express mineralocorticoid receptors and the enzyme necessary for
399 mineralocorticoid-specific action, known as 11- β hydroxysteroid dehydrogenase 2
400 (Geerling et al., 2006; Geerling and Loewy, 2006). In contrast, humoral AngII acts initially
401 on AngII type 1 (AT1) receptors in circumventricular organs, such as the subfornical organ

402 and the organ vasculosum of the lateral terminalis (Lind et al., 1985; McKinley et al.,
403 1992b; Tanaka et al., 1986; Weiss and Hatton, 1990). Mineralocorticoids exert minor up-
404 regulation of AT1 receptor levels and AT1 signaling (Grafe et al., 2014; Shelat et al.,
405 1999a; Shelat et al., 1998; Shelat et al., 1999b). As a parallel mechanism,
406 mineralocorticoids may exert a cooperative behavioral effect with AngII by acting
407 elsewhere in a broader circuit. For example, mineralocorticoids appear to remove an
408 inhibitory influence on sodium appetite, namely oxytocin activity in the PVN (Blackburn et
409 al., 1992; Grafe et al., 2014; Roesch et al., 2001; Stricker and Verbalis, 1987).

410 A behaviorally relevant downstream connection of both mineralocorticoid- and
411 AngII-sensing brain regions is the lateral hypothalamus (Camacho and Phillips, 1981),
412 which is essential for the normal expression of sodium appetite (Dayawansa et al., 2011).
413 The lateral hypothalamus, in turn, projects to the mesolimbic dopamine system via the
414 medial forebrain bundle (Berk and Finkelstein, 1981; Saper et al., 1979). Relevant
415 projections from the lateral hypothalamus to the VTA may include orexin neurons (Fadel
416 and Deutch, 2002; Narita et al., 2006; Peyron et al., 1998), which promote incentive-
417 based effort in the VTA (Borgland et al., 2009; Choi et al., 2010). Furthermore, stimulation
418 of the lateral hypothalamus promotes sodium appetite (Liedtke et al., 2011). Likewise,
419 AngII treatment enhances the release of striatal dopamine in freely moving rats within
420 minutes, which then is associated with increased drinking behavior (Brown et al., 1996;
421 Hoebel et al., 1994). Thus, a possible link between mineralocorticoid and AngII-sensing
422 brain regions and midbrain dopamine neurons may be lateral hypothalamic orexin
423 neurons. Future studies are needed to illuminate the details of this broader circuit.

424

425 *Sodium appetite and motivation systems*

426 The mesolimbic pathway is part of a motor loop that links the incentive properties
427 of stimuli with the effort exerted to approach or avoid them (Berridge, 2007; Salamone et
428 al., 2007). Neural adaptations in this brain region are associated with motivated
429 behaviors (Gu et al., 2010). AngII-induced cFos expression in the mesolimbic area is
430 likely the result of transynaptic activation because the VTA does not have AngII receptors
431 (Song et al., 1992). Previous studies documented AngII-induced activation of the
432 hypothalamus and taste nuclei (Han and Rowland, 1995; Houtp et al., 1998; McKinley et
433 al., 1992a; Thunhorst et al., 1998; Vivas et al., 1995), but this is the first report of cFos
434 activation in the VTA.

435 The VTA includes many dopaminergic neurons (Oades and Halliday, 1987). The
436 AngII-induced increase in tyrosine hydroxylase levels supports the notion that dopamine
437 neurons are activated by AngII treatment. A similarly rapid increase in tyrosine
438 hydroxylase levels occurs during exposure to cocaine-paired contexts (Liang et al., 2012).
439 AngII-induced cFos expression may contribute to the up-regulation of tyrosine
440 hydroxylase (Gheea et al., 1998). DOC treatment markedly increased the
441 phosphorylation of tyrosine hydroxylase, which augments catalytic activity of this rate-
442 limiting enzyme (Dunkley et al., 2004). The combined actions of AngII and DOC on
443 neuronal activation and tyrosine hydroxylase would be expected to produce a substantial
444 increase in dopamine release at axonal targets. Future studies are needed to examine
445 the consequences of these hormonal actions at downstream nodes in the motivation
446 circuit.

447 Regarding the ventral striatum, neurons there exhibit neural plasticity during
448 various motivational states, including depletion-induced sodium appetite (Roitman et al.,
449 2002). As in the VTA, AngII treatment increased cFos levels without a detectable
450 augmentation by DOC pretreatment. Unlike the VTA, all hormone treatments increased
451 phosphorylation of tyrosine hydroxylase levels with detectable increases in total levels of
452 this enzyme. It will be important for future studies to investigate the neurophysiological
453 consequences and potential rewiring that may occur in the accumbens in response to
454 AngII- versus DOC/AngII-induced changes in this dopamine pathway.

455

456 *Conclusions*

457 In summary, the DOC/AngII-induced sodium appetite involves a shift in the effort
458 value of sodium versus water consumption, noted in the break point for lever pressing.
459 DOC and AngII affect the mesolimbic dopamine pathway differently, with AngII treatment
460 increasing cFos activation in the VTA and nucleus accumbens, and DOC increasing
461 phosphorylation of tyrosine hydroxylase. Thus, these two hormones may act in parallel
462 to enhance mesolimbic dopamine transmission through complementary mechanisms.

463

464

465 **Acknowledgements**

466

467 This work was supported by the National Institutes of Health Grants R01 HL091314. The
468 advice and expertise of Dr. Scott E. Kanoski was indispensable for the progressive ratio
469 experiment. We thank Dr. Harvey J. Grill for the use of his progressive ratio conditioning
470 equipment. We appreciate the assistance of Olivia Kim and Alison Krieger in some of the
471 data analysis.

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493 **References**

494

495 Andersson, B., 1977. Regulation of body fluids. *Annu.Rev.Physiol.* 39, 185-200.

496 Barbano, M.F., Cador, M., 2006. Differential regulation of the consummatory, moti-
497 vational, and anticipatory aspects of feeding behavior by dopaminergic and
498 opiodergic drugs. *Neuropsychopharmacology* 31, 1371–1381.

499 Berk, M.L., Finkelstein, J.A., 1981. Afferent projections to the preoptic area and
500 hypothalamic regions in the rat brain. *Neuroscience* 6, 1601-1624.

501 Berridge, K.C., 2007. The debate over dopamine's role in reward: the case for incentive
502 salience. *Psychopharmacology* 191, 391-431.

503 Berridge, K.C., Flynn, F.W., Schulkin, J., Grill, H.G., 1984. Sodium depletion enhances
504 salt palatability in rats. *Behav Neurosci* 98, 652-660.

505 Blackburn, R.E., Demko, A.D., Hoffman, G.E., Stricker, E.M., Verbalis, J.G., 1992. Central
506 oxytocin inhibition of angiotensin-induced salt appetite in rats. *Am. J. Physiol.* 263,
507 R1347-R1353.

508 Borgland, S.L., Chang, S.-J., Bowers, M.S., Thompson, J.L., Vittoz, N., Floresco, S.B.,
509 Chou, J., Chen, B.T., Bonci, A., 2009. Orexin A/hypocretin-1 selectively promotes
510 motivation for positive reinforcers. *J. Neurosci.* 29, 11215-11225.

511 Brown, D.C., Steward, L.J., Ge, J., Barnes, N.M., 1996. Ability of angiotensin II to
512 modulate striatal dopamine release via the AT1 receptor in vitro and in vivo. *Br.*
513 *J. Pharmacol.* 118:, 414-420.

514 Buggy, J., Jonklaas, J., 1984. Sodium appetite decreased by central angiotensin
515 blockade. *Physiol.Behav.* 32, 737-742.

516 Camacho, A., Phillips, M.I., 1981. Horseradish peroxidase study in rat of the neural
517 connections of the organum vasculosum of the lamina terminalis. *Neuroscience*
518 *Letters* 25 201-204.

519 Carelli, R.M., 2002. The nucleus accumbens and reward: neurophysiological
520 investigations in behaving animals. *Behav. Cogn. Neurosci. Rev.* 1, 281-296

521 Choi, D.L., Davis, J.F., Fitzgerald, M.E., Benoit, S.C., 2010. The role of orexin-A in food
522 motivation, reward-based feeding behavior and food-induced neuronal activation
523 in rats. *Neurosci.* 167, 11-20.

524 Clark, J.J., Bernstein, I.L., 2006. Sensitization of salt appetite is associated with increased
525 "wanting" but not "liking" of a salt reward in the sodium-deplete rat. *Behav.*
526 *Neurosci.* 120, 206-210.

527 Conover, K.L., Woodside, B., Shizgal, P., 1994. Effects of sodium depletion on
528 competition and summation between rewarding effects of salt and lateral
529 hypothalamic stimulation in the rat. *Behav. Neurosci.* 108, 549-558.

530 Davis, J.F., Choi, D.L., Schurdak, J.D., Fitzgerald, M.F., Clegg, D.J., Lipton, J.W.,
531 Figlewicz, D., Benoit, S.C., 2011. Leptin regulates energy balance and motivation
532 through action at distinct neural circuits. *Biol.Psychiatry* 69, 668-674.

533 Dayawansa, S., Peckins, S., Ruch, S., Norgren, R., 2011. Parabrachial and hypothalamic
534 interaction in sodium appetite. *Am. J. Physiol.* 300, R1091-R1099.

535 Dunkley, P.R., Bobrovskaya, L., Graham, M.E., Von Nagy-Felsobuki, E.I., Dickson, P.W.,
536 2004. Tyrosine hydroxylase phosphorylation: regulation and consequences. *J.*
537 *Neurochem.* 91, 1025–1043.

538 Epstein, A.N., Fitzsimons, J.T., Simons, B.J., 1969. Drinking caused by the intracranial
539 injection of angiotensin into the rat. *J. Physiol.* 200, 98P-100P.

540 Fadel, J., Deutch, A.Y., 2002. Anatomical substrates of orexin-dopamine interactions:
541 lateral hypothalamic projections to the ventral tegmental area. *Neuroscience*
542 *Letters* 111, 379-387.

543 Floresco, S.B., St. Onge, J.R., Ghods, S., S., Winstanley, S.A., 2008. Cortico-limbic-
544 striatal circuits subserving different forms of cost-benefit decision making. *Cogn.*
545 *Affect. Behav. Neurosci.* 8, 375-389.

546 Fluharty, S.J., Epstein, A.N., 1983. Sodium appetite elicited by intracerebroventricular
547 infusion of angiotensin II in the rat. II. synergistic interaction with systemic
548 mineralocorticoids. *Behav Neurosci* 97, 746-758.

549 Fluharty, S.J., Sakai, R.R., 1995. Behavioral and cellular analysis of adrenal steroid and
550 angiotensin interactions mediating salt appetite, in: Fluharty, S.J., Morrison, A.R.,
551 Sprague, J., Stellar, E. (Eds.), *Prog. Psychobiol. Physiol. Psych.* Academic Press,
552 San Diego, CA, pp. 177-212.

553 Garcia, J.M., Curtis, K.S., Contreras, R.J., 2008. Behavioral and electrophysiological
554 taste responses change after brief or prolonged dietary sodium deprivation. *Am. J.*
555 *Physiol.* 295, R1754-R1761.

556 Geerling, J.C., Kawata, M., Loewy, A.D., 2006. Aldosterone-sensitive neurons in the rat
557 central nervous system. *J. Comp. Neurol.* 494, 515-527.

558 Geerling, J.C., Loewy, A.D., 2006. Aldosterone-sensitive neurons in the nucleus of the
559 solitary tract: bidirectional connections with the central nucleus of the amygdala. *J*
560 *Comp Neurol* 497, 646–657.

561 Gheea, M., Bakerb, H., Millerc, J.C., Ziff, E.B., 1998. AP-1, CREB and CBP transcription
562 factors differentially regulate the tyrosine hydroxylase gene. *Mol. Brain Res.* 55,
563 101–114.

564 Grafe, L.A., Takacs, A.E., Yee, D.K., Flanagan-Cato, L.M., 2014. The role of the
565 hypothalamic paraventricular nucleus and the organum vasculosum lateral
566 terminalis in the control of sodium appetite in male rats. *J. Neurosci.* 34, 9249-
567 9260.

568 Grondin, M.E., Gobeil-Simard, A., Drolet, G., Mougnot, D., 2011. Na⁺ appetite induced
569 by depleting extracellular fluid volume activates the enkephalin/mu-opioid receptor
570 system in the rat forebrain *Neuroscience Letters* 192, 398-412,.

571 Gu, H., Salmeron, B.J., Ross, T.J., Geng, X., Zhan, W., Stein, E.A., Yang, Y., 2010.
572 Mesocorticolimbic circuits are impaired in chronic cocaine users as demonstrated
573 by resting-state functional connectivity. *Neuroimage* 53, 593-601.

574 Han, L., Rowland, N.E., 1995. Sodium depletion and Fos-immunoreactivity in lamina
575 terminalis. *Neurosci. Lett.* 193, 173-176.

576 Hodos, W., 1961. Progressive ratio as a measure of reward strength. *Science* 134 943-
577 944.

578 Hoebel, B.G., Rada, P., Mark, G.P., Hernandez, L., 1994. The power of integrative
579 peptides to reinforce behavior by releasing dopamine. *Ann. N.Y. Acad. Sci.* 739,
580 36-41.

581 Houpt, T.A., Smith, G.P., Joh, T.H., Frankmann, S.P., 1998. c-fos-like immunoreactivity
582 in the subfornical organ and nucleus of the solitary tract following salt inake by
583 sodium depleted rats. *Physiol. Behav.* 63, 505-510.

584 Johnson, A.K., Thunhorst, R.L., 1997. The neuroendocrinology of thirst and salt appetite:
585 visceral sensory signals and mechanisms of central integration. *Front.*
586 *Neuroendocrinol.* 18, 292-353.

587 Kelley, A.E., Baldo, B.A., Pratt, W.E., Will, M.J., 2005. Corticostriatal-hypothalamic
588 circuitry and food motivation: integration of energy, action and reward. *Physiol.*
589 *Behav.* 86 773–795.

590 Kovács, K.J., 2008. Measurement of immediate-early gene activation- c-fos and beyond.
591 *J. Neuroendocrinol.* 20, 665-672.

592 Kraulis, I., Foldes, G., Traikov, H., Dubrovsky, B., Birmingham, M.K., 1975 Distribution,
593 metabolism and biological activity of deoxycorticosterone in the central nervous
594 system. *Brain Res.* 88, 1-14.

595 Liang, J., Ma, S.S., Li, Y.J., Ping, X.J., Hu, L., Cui, C.L., 2012. Dynamic changes of
596 tyrosine hydroxylase and dopamine concentrations in the ventral tegmental area-
597 nucleus accumbens projection during the expression of morphine-induced
598 conditioned place preference in rats. . *Neurochem. Res.* 37, 1482-1489.

599 Liedtke, W.B., McKinley, M.J., Walker, L.L., Zhang, H., Pfenning, A.R., Drago, J.,
600 Hochendoner, S.J., Hilton, D.L., Lawrence, A.J., Denton, D.A., 2011. Relation of
601 addiction genes to hypothalamic gene changes subserving genesis and
602 gratification of a classic instinct, sodium appetite. *Proc. Natl. Acad. Sci. U.S.A.* 108,
603 12509-12514.

604 Lind, R.W., Swanson, L.W., Sawchenko, P.E., 1985. Anatomical evidence that neural
605 circuits related to the subfornical organ contain angiotensin II. *Brain Res Bull* 15,
606 79-82.

607 Lucas, L.R., Grillo, C.A., McEwen, B.S., 2003. Involvement of mesolimbic structures in
608 short-term sodium depletion: in situ hybridization and ligand-binding analyses.
609 *Neuroendocrinology* 77, 406-415.

610 Lundy, R.F., Jr., 2008. Gustatory hedonic value: Potential function for forebrain control of
611 brainstem taste processing. *Neurosci. Biobehav. Rev.* 32, 1601-1606.

612 McKinley, M.J., Badoer, E., Oldfield, B.J., 1992a. Intravenous angiotensin II induces fos-
613 immunoreactivity in circumventricular organs of the lamina terminalis. *Brain Res.*
614 594, 295-300.

615 McKinley, M.J., Bicknell, R.J., Hards, D., McAllen, R.M., Vivas, L., Weisinger, R.S.,
616 Oldfield, B.J., 1992b. Efferent neural pathways of the lamina terminalis subserving
617 osmoregulation. *Prog. Brain Res.* 91, 395-402.

618 Miller, R.L., Loewy, A.D., 2014. 5-HT neurons of the area postrema become c-Fos-
619 activated after increases in plasma sodium levels and transmit interoceptive
620 information to the nucleus accumbens. *Am. J. Physiol.* 306, R663-R673.

621 Narita, M., Nagumo, Y., Hashimoto, S., Narita, M., Khotib, J., Miyatake, M., Sakurai, T.,
622 Yanagisawa, M., Nakamachi, T., Shioda, S., Suzuki, T., 2006. Direct involvement
623 of orexinergic systems in the activation of the mesolimbic dopamine pathway and
624 related behaviors induced by morphine. *J. Neurosci.* 26, 398-405.

625 Oades, R.D., Halliday, G.M., 1987. Ventral tegmental (A10) system: neurobiology. 1.
626 Anatomy and connectivity. *Brain Res.* 434, 117-165.

627 Peyron, C., Tighe, D.K., van den Pol, A.N., de Lecea, L., Heller, H.C., Sutcliffe, J.G.,
628 Kilduff, T.S., 1998. Neurons containing hypocretin (orexin) project to multiple
629 neuronal systems. *J. Neurosci.* 18, 9996-10015.

630 Phillips, P.E., Walton, M.E., Jhou, T.C., 2007. Calculating utility: preclinical evidence for
631 cost-benefit analysis by mesolimbic dopamine. *Psychopharmacology (Berl)* 191,
632 483–495.

633 Roesch, D.M., Blackburn-Munro, R.E., Verbalis, J.G., 2001. Mineralocorticoid treatment
634 attenuates activation of oxytocinergic and vasopressinergic neurons by icv ANG
635 II. *Am. J. Physiol* 280, R1853-R1864.

636 Roitman, M.F., Na, E., Anderson, G., Jones, T.A., Bernstein, I.L., 2002. Induction of a salt
637 appetite alters dendritic morphology in nucleus accumbens and sensitizes rats to
638 amphetamine. *J. Neurosci.* 22, RC225 (221-225).

639 Roitman, M.F., Patterson, T.A., Sakai, R.R., Bernstein, I.L., Figlewicz, D.L., 1999. Sodium
640 depletion and aldosterone decrease dopamine transporter activity in rat nucleus
641 accumbens but not striatum. *Am. J. Physiol.* 276, R1339-R1345.

642 Sakai, R.R., Nicolaidis, S., Epstein, A.N., 1986. Salt appetite is suppressed by
643 interference with angiotensin II and aldosterone. *Am. J. Physiol* 251, R762-R768.

644 Salamone, J.D., Correa, M., Farrar, A., Mingote, S.M., 2007. Effort-related functions of
645 nucleus accumbens dopamine and associated forebrain circuits.
646 *Psychopharmacology* 191, 461-482.

647 Salamone, J.D., Correa, M., Farrar, A.M., Nunes, E.J., Pardo, M., 2009. Dopamine,
648 behavioral economics, and effort. *Front. Behav. Neurosci.* 3, 13.

649 Saper, C.B., Swanson, L.W., Cowan, W.M., 1979. An autographic study of the efferent
650 connections of the lateral hypothalamic area in the rat. *J. Comp. Neurology* 183,
651 689-706.

652 Sclafani, A., Ackroff, K., 2003. Reinforcement value of sucrose measured by progressive
653 ratio operant licking in the rat. *Physiol. Behav.* 79, 663-670.

654 Shade, R.E., Blair-West, J.R., Carey, K.D., Madden, L.J., Weisinger, R.S., Denton, D.A.,
655 2002. Synergy between angiotensin and aldosterone in evoking sodium appetite
656 in baboons. *Am. J. Physiol* 283, R1070-R1078.

657 Shekhtman, E., Geerling, J.C., Loewy, A.D., 2007. Aldosterone-sensitive neurons of the
658 nucleus of the solitary tract: multisynaptic pathway to the nucleus accumbens *J.*
659 *Comp. Neurol.* 501, 274-289.

660 Shelat, S.G., Flanagan-Cato, L.M., Fluharty, S.J., 1999a. Glucocorticoid and
661 mineralocorticoid regulation of angiotensin II type 1 receptor binding and inositol-
662 trisphosphate signaling in WB cells *J. Endocrinology* 162, 381-391.

663 Shelat, S.G., Fluharty, S.J., Flanagan-Cato, L.M., 1998. Adrenal steroid regulation of
664 central angiotensin II receptor subtypes and oxytocin receptors in rat brain. *Brain*
665 *Research* 807, 135-146.

666 Shelat, S.G., King, J.L., Flanagan-Cato, L.M., Fluharty, S.J., 1999b. Mineralocorticoid and
667 glucocorticoid interactions on salt appetite and angiotensin II receptor binding in
668 rat brain and pituitary. *Neuroendocrinology* 69, 339-351.

669 Song, K., Allen, A.M., Paxinos, G., Mendelsohn, F.A.O., 1992. Mapping of angiotensin II
670 receptor subtype heterogeneity in rat brain. *J. Comp. Neurol.* 316, 467-484.

671 Starr, L.J., Rowland, N.E., 2006. Characteristics of salt appetite in chronically sodium-
672 depleted rats using a progressive ratio schedule of procurement. *Physiol. Behav.*
673 88, 433-442.

674 Stricker, E.M., Verbalis, J.G., 1987. Central inhibitory control of sodium appetite in rats:
675 correlation with pituitary oxytocin secretion. *Behav. Neurosci.* 101, 560-567.

676 Tanaka, J., Kaba, H., Saito, H., Seto, K., 1986. Subfornical organ efferents influence the
677 activity of median preoptic neurons projecting to the hypothalamic paraventricular
678 nucleus in the rat. *Exp. Neurol.* 93, 647-651.

679 Tekin, I., Roskoski, R., Jr., Carkaci-Salli, N., Vrana, K.E., 2014. Complex molecular
680 regulation of tyrosine hydroxylase. *J. Neural Transmission* in press.

681 Thunhorst, R.L., Xu, Z., Cicha, M.Z., Zardetto-Smith, A.M., Johnson, A.K., 1998. Fos
682 expression in rat brain during depletion-induced thirst and sodium appetite. *Am. J.*
683 *Physiol.* 274, R1807-R1814.

684 Tindell, A.J., Smith, K.S., Pecina, S., Berridge, K.C., Aldridge, J.W., 2006. Ventral
685 pallidum firing codes hedonic reward: when a bad taste turns good. *J.*
686 *Neurophysiol.* 96, 2399-2409.

687 Vivas, L., Pastuskovas, C.V., Tonelli, L., 1995. Sodium depletion induces Fos
688 immunoreactivity in the circumventricular organs of the lamina terminalis. *Brain*
689 *Res.* 679.

690 Weiss, M.L., Hatton, G.I., 1990. Collateral input to the paraventricular and supraoptic
691 nuclei in rat. I. Afferents from the subfornical organ and the anteroventral third
692 ventricle region. *Brain Res. Bull.* 24, 231-238.

693 Zhang, D.M., Stellar, E., Epstein, A.N., 1984. Together intracranial angiotensin and
694 systemic mineralocorticoid produce avidity for salt in the rat. *Physiol. Behav.* 32,
695 677-681.

696

697 **Figure Legends**

698

699 **Figure 1.** Drawings of rat brain coronal sections depicting the regions of interest for both
700 cFos immunohistochemistry (**Panel A**) and tyrosine hydroxylase activation assays (**Panel**
701 **B**). **Panel A.** Boxes surrounding each brain region represent the areas analyzed for
702 cFos cell counts. **Panel B.** Circles within each brain region represent 0.75-mm
703 micropunches collected for the tyrosine hydroxylase western blots. The brain regions
704 examined include both the core and shell of the accumbens, as well as the VTA.
705 Abbreviations: AcbC = Core of the accumbens, AcbS = Shell of the accumbens, CPu=
706 caudate putamen, ml = medial lemniscus, LS = lateral septal nucleus, PAG =
707 periaqueductal gray, VTA= ventral tegmental area.

708

709 **Figure 2.** DOC/AngII treatment increases relative motivation for sodium (n = 5-6/group).
710 **Panel A.** Line graphs illustrating cumulative presses over time for 3% Saline and Water
711 after Vehicle, DOC, AngII, or DOC plus AngII treatments. **Panel B.** Bar graphs illustrating
712 total lever presses for 3% saline and water after Vehicle, DOC, AngII, or DOC plus AngII
713 treatments. AngII-only treatment increased bar presses for water and sodium, whereas
714 DOC plus AngII treatment mainly increased presses for sodium. **Panel C.** Bar graphs
715 illustrating the breakpoint ratio of sodium to water after Vehicle, DOC, AngII, or DOC plus
716 AngII. DOC plus AngII causes the highest sodium to water breakpoint ratio.
717 Abbreviations: AngII= Angiotensin II, DOC = deoxycorticosterone acetate, Veh = vehicle.

718

719 **Figure 3.** DOC/AngII treatment increases cFos expression in the ventral tegmental area
720 and the accumbens (n = 3-6 per group). Bar graphs illustrate the cFos cell counts in the
721 ventral tegmental area and accumbens after either vehicle or DOC pretreatment followed
722 by *icv* vehicle or AngII. Representative images of the VTA and accumbens (coronal
723 plane, 10x) in each treatment condition are shown above the bar graphs. AngII and
724 DOC/AngII treatment increased cFos expression in the VTA, with DOC/AngII inducing the
725 most immunostaining. In the core and shell of the nucleus accumbens, each hormone
726 treatment induced cFos immunostaining compared with vehicle; DOC/AngII induced the
727 highest amount of cFos immunostaining. Asterisks indicate a significant increase
728 compared with the Veh/Veh group. Abbreviations: AngII = Angiotensin II, DOC =
729 deoxycorticosterone acetate, NuAcc= Accumbens, Veh = Vehicle, VTA= Ventral
730 Tegmental Area

731

732 **Figure 4.** DOC treatment increases tyrosine hydroxylase activation in the VTA and the
733 accumbens (n = 3/group). **Panel A.** Bar graphs illustrate tyrosine hydroxylase levels in
734 the VTA and the accumbens (core and shell) after oil versus DOC pretreatment and *icv*
735 saline versus AngII. Representative western blot images of tyrosine hydroxylase are
736 shown above each quantified bar. Each treatment increased tyrosine hydroxylase
737 expression in the VTA. **Panel B.** Bar graphs illustrate phosphorylated tyrosine
738 hydroxylase levels in the VTA and the accumbens (core and shell) after either oil or DOC
739 pretreatment followed by *icv* treatments with AngII. Only DOC pretreatment significantly
740 increased phosphorylated tyrosine hydroxylase expression compared to vehicle in the
741 VTA and accumbens core. However, each treatment increased phosphorylated tyrosine

742 hydroxylase expression in the accumbens shell. Panel C. Representative western blot
743 images labeled with corresponding treatment groups and brain regions assayed for
744 tyrosine hydroxylase (left) and phosphorylated tyrosine hydroxylase (right).
745 Abbreviations: AngII = Angiotensin II, Acc= Accumbens, DOC = deoxycorticosterone
746 acetate phospho-TH = phosphorylated tyrosine hydroxylase, TH = tyrosine hydroxylase,
747 Veh = Vehicle, VTA= ventral tegmental area. Asterisks indicate $p < 0.05$ compared with
748 the Veh/Veh group.

749

750 **Figure 5.** Diagram summarizing the effects of DOC and AngII on the mesolimbic system
751 and behavior. Abbreviations: AngII = Angiotensin II, NuAcc= Nucleus Accumbens, DOC
752 = deoxycorticosterone acetate, p-TH = phosphorylated tyrosine hydroxylase, TH =
753 tyrosine hydroxylase, VTA= ventral tegmental area.

754

755

756

757

758

759

760

761